

PRINCIPLES AND PROSPECTS FOR  
BIOREMEDIATION OF PCBs IN SOILS AND  
SEDIMENTS



James M. TIEDJE  
Distinguished Professor and  
Director  
Center For Microbial Ecology  
Michigan State University  
tiedjej@msu.edu




### **Current Biological Principles For Bioremediation Of PCBs**

- **Highly chlorinated PCBs (Aroclors) can be reduced to a lesser chlorinated species by anaerobic dehalogenating communities**
- **The lesser chlorinated PCB species can be oxidized by aerobic biphenyl degraders yielding chlorinated benzoates, pentadienes and HOPDAs**
- **Chlorobenzoates can be mineralized by yet another group of bacteria**
- **These three independently occurring processes can be combined in a singular two-phase anaerobic-aerobic remediation scheme**


## Rationale for anaerobic/aerobic biotreatment

Less aerobically degradable

More aerobically degradable




# Chromatographic Profiles - Pattern



## Congener Specificity

Dechlorination Activity	Susceptible Chlorines
M	Flanked & unflanked <i>meta</i>
Q	Flanked & unflanked <i>para</i>
H	Flanked <i>para</i> Doubly flanked <i>meta</i>
H'	Flanked <i>para</i> <i>Meta</i> of 2,3- & 2,3,4- groups
P	Flanked <i>para</i>
N	Flanked <i>meta</i>
T	<i>Meta</i> of hepta- & octa-CBs
LP	Unflanked <i>para</i>

## Evaluation of *in situ* PCB Dechlorination




## Occurrence of PCB Dechlorination

- Acushnet Estuary (MA)
- Hudson River
- Industrial lagoons
- River Raisin (MI)
- Sheboygan River (WI)
- Silver Lake (MA)
- Escambia Bay (FL)
- Hoosic River (MA)
- Lake Ketelmeer/Rhine River (Netherlands)
- Waukegan Harbor (IL)
- Wood's Pond (MA)




Observed extent of dechlorination tends to taper off below 50 to 100 ppm.

### FeSO<sub>4</sub> Amendment Extent of Dechlorination





## Effect of $\text{FeSO}_4$ on Aroclor 1242 Dechlorination by Hudson River Microorganisms



No amendment:

Process M (*meta*) dechlorination

25% of Cl removed

*Ortho* & *para* substituted products:

2-CB, 2,2'-CB, 2,6-CB, 2,4'-CB,

2,2',4-CB, 2,4,4'-CB

With  $\text{FeSO}_4$ :


Adds process Q (*para*) dechlorination

50% of Cl removed

*Ortho* substituted products:

2-CB, 2,2'-CB, 2,6-CB

## Sequential Inoculations Extent of Dechlorination




## Importance of PCB Dechlorination

- Is an intrinsic process
- Has potential for bioremediation
- Products are more aerobically degradable
- Products are generally less toxic
- But, degradation is usually incomplete with especially *ortho*-PCBs remaining, hence, an aerobic phase is needed



## Barriers to Aerobic Biodegradation

- Limited to lesser chlorinated congeners
- Co-metabolic
  - Requires induction
  - Yields no growth
- Incomplete - accumulates potentially problematic compounds

## Recombinant PCB degradation Pathways




13




**Differential expression of dehalogenases in response to CBAs.**  
Reverse transcription (RT)-PCR of total RNA from *Nocardioides* sp. KZ4N.

- ◆ Halohydroperoxidase/catalase (*catA*) to use for upgrading pathway for *ortho* + *para*-PCBs
- ◆ The *catA* is related to (bromo) hydroxyperoxidases for anoxic chlorination/dechlorination of chloramphenicol
- ◆ The laboratory evolution to target enzyme(s) for reductive dechlorination of both CBAs and PCBs
- ◆ The *catA* family members to use for the combinatorial mutagenesis




### Schematic Representation of the Sediment Experiment









### Molecular Tracking of RHA 1 (fcbB) in Picatinny Arsenal Soil Using Real Time PCR



**Population dynamics of RHA1(*fc*b) and LB400(*oh*b) during aerobic treatment of Aroclor 1242 anaerobic dechlorination products (Red Cedar River sediment)**



Anaerobic/Aerobic Degradation of Aroclor 1242



Laboratory scale two-phase PCB remediation  
(Red Cedar River sediment)

- Enhanced anaerobic dechlorination of Aroclor resulted in shifting from highly to lower chlorinated congeners
- Bioaugmentation with engineered aerobic G+/G- GEM cassette LB400(*ohb*)+RHA1(*fc*b) resulted in efficient degradation of the remaining PCBs

## PCB-growing Dechlorinating GEM Advantages

- ◆ **Increased biomass**  
smaller inoculum
- ◆ **Reduced HOPDA**  
diminished toxicity
- ◆ **Issues remaining: Still incomplete mineralization of higher chlorinated PCB species**  
lower degradation rates, incomplete Cl-HOPDA removal

### Effect of the *ohb* (pRO41) and *pcb*(pRHD34) genes on PCB degradation


Strain(plasmid)/ inducer	% PCB degraded	Mix M, nom. 1mM			Mix C, nom. 1 mM			
		OD <sub>600</sub>	Cl <sup>-</sup> Release, mM	HOPDA $\lambda=394$	% PCB degraded	OD <sub>600</sub>	Cl <sup>-</sup> Release, mM	HOPDA $\lambda=394$
LB400(pRT1) <sub>Bph</sub>	83.8	<b>0.19</b>	1.33	<b>0.476</b>	93.7	<b>0.21</b>	0.91	<b>0.066</b>
LB400(pRO41) <sub>25CBA</sub>	75	<b>0.24</b>	1.98	<b>0.164</b>	93.2	<b>0.35</b>	1.47	<b>Not detected</b>
LB400(pRO41) <sub>25CBA</sub> + RHA1(pRHD34) <sub>4CBA</sub>	67.6	0.19	1.5	<b>0.139</b>	91.3	<b>0.30</b>	1.42	<b>0.069</b>

20

# Biological Activity

## Dechlorination of Congeners with AhR-Mediated Activity

### Aroclor 1242



## AhR Mediated Toxicity Reduction

Sample	TEQ <sub>assay</sub> *	TEQ <sub>calc</sub> *
Non-dechlorinated 1242	3.1	5.7
SL dechlorinated 1242	<6 X 10 <sup>-2</sup> **	7.8 X 10 <sup>-2</sup>
RR dechlorinated 1242	<6 X 10 <sup>-2</sup> **	4.7 X 10 <sup>-2</sup>

\*Units are pmole TEQ/Σ μmoles of PCBs.

\*\*TEQ<sub>assay</sub> was less than the MDL.

## Biological Activity Summary

Biological Activity	Anaerobic Dechlorination	Aerobic Di-OHs
AhR mediated	Reduced/eliminated	Inactive
AP-transcription	Reduced/eliminated	Inactive
Fertilization	Reduced/eliminated	-----
Neutrophil activation	Unaffected	Inactive
Insulin release	Reduced/eliminated	Decreased
Estrogenicity	-----	Inactive
Uterine contractility	Increased	-----



## Genome of *Burkholderia* Strain LB400

- Best known PCB degrader sequenced
- 9.5 Megabases, largest bacterial genome yet sequenced
- Has multiple chromosomes
- Has many aromatic degradation genes



## Differential Expression by a PCB Degradar

4,000 genes of  
*Burkholderia* strain LB400



Genes (mRNA) expressed during growth on succinate (green, Cy3) vs. genes expressed during growth on rich media (red, Cy5), and expressed under both conditions (yellow)

## Hudson River PCBs: Lessons from Dated Sediment Cores

Richard Bopp and Edward Shuster, Rensselaer Polytechnic Institute

Steven Chillrud, Lamont Doherty Earth Observatory of Columbia University

Frank Estabrooks, New York State Department of Environmental Conservation


NIEHS Superfund Basic Research Program

Grant to Mount Sinai Medical Center

Philip J. Landrigan, P.I.

OCTOBER 2002

**26**









## Basics of Our Radionuclide Dating of Sediment Cores

- Cs-137 - supplied by global fallout from atmospheric testing of nuclear weapons. In “ideal” sediment cores the Cs-137 profile provides at least two time horizons - a “first appearance” in about 1954 and a maximum activity in 1963-64.


- Be-7 - a short-lived natural radionuclide (half life = 53 days). Detection of Be-7 in near surface sections of sediment cores confirms the presence of a significant component of particles deposited within about a year of core collection.





“Well-dated” cores have been collected from the Hudson at mile points 203, 193.8, 188.5, 177.6, 163.6, 152.7, 91.8, 88.6, 59, 54, 43, 21.6, 3.0, and -1.7. Samples of all core sections have been archived.






Despite major “dilution” with cleaner particles from tributaries, the early 1970s peak in PCB levels can be traced downstream to the NY/NJ Harbor.

The best evidence for continuing transport of PCBs from sediments of the Thompson Island Pool was provided by weekly water column monitoring.

Thompson Island Dam:  
TID-WEST and TID-PRW2  
with Fort Edward





In situ reductive dechlorination of PCBs has been studied using paired samples from co-located sediment cores.


- This process results in dramatic alteration of PCB composition in upper Hudson sediments, but has not progressed as far as lab studies would suggest.


- In sediments from the freshwater reach of the tidal Hudson, reductive dechlorination can be easily detected, but the overall effect on PCB composition has been minor.

Matching contaminant profiles in paired, co-located cores












“Other” significant contaminants in upper Hudson sediments include Pb, Cd, Cr, Zn, and dioxins.










The importance of other PCB sources can be determined from analyses of dated sediments from “indicator” sites. Site 9 is Newtown Creek, a municipal wastewater indicator. Sites 6 & 7 (NYC drinking water reservoirs) and Central Park Lake are atmospheric deposition indicators.


**Table 1.** Contaminant levels in sediment samples deposited in the mid 1980s.

	Newtown Creek site 9	NY/NJ harbor site 8	Kingston site 5
Organic matter (%) <sup>a</sup>	12.8	9.5 (64%) <sup>b</sup>	3.6
pp'-DDD (ppb)	62	35 (52%)	6.2
γ-chlordane (ppb)	28	16 (57%)	<1
total PCBS (ppm)	1.91	1.47 (61%)	0.79
copper (ppm)	604	207 (31%)	31
lead (ppm)	466	223 (41%)	53
zinc (ppm)	748	358 (35%)	145
cadmium (ppm)	10.5	4.03 (31%)	1.12

<sup>a</sup>Based on weight loss on ignition at 375°C overnight.

<sup>b</sup>Values in parentheses represent the sewage contribution to contaminant levels in this sample calculated as ((NY/NJ harbor – Kingston)/ (Newtown Creek – Kingston)) x 100.





Conclusions:

- Analyses of sections from dated sediment cores is the single most efficient means of understanding the sources, distribution, and fate of particle-associated contaminants in natural water systems. (So I'm biased.)

- A multi-contaminant/tracer approach provides critical support for any interpretation of the behavior of any specific particle-associated contaminant (PCBs) in a natural water system (the Hudson).

↳ Recent significant developments in the Hudson River PCB story include stable lead isotopes as tracers of upper Hudson particles; plutonium isotopes as tracers of Mohawk River particles; and stable isotopes as tracers of reductive dechlorination of PCBs.

# Thank You

After viewing the links to additional resources,  
please complete our online feedback form.

**Thank You**

[Links to Additional Resources](#)

