

# Benthic Flux of Dissolved Nickel into the Water Column of South San Francisco Bay

By Brent R. Topping, James S. Kuwabara, Francis Parchaso, Stephen W. Hager, Andrew J. Arnsberg, and Fred Murphy



Open File Report 01-89

Core Retrieval photo by Oscar Mace

U.S. Department of Interior  
U.S. Geological Survey

# **Benthic Flux of Dissolved Nickel into the Water Column of South San Francisco Bay**

By Brent R. Topping<sup>1</sup>, James S. Kuwabara<sup>2</sup>, Francis Parchaso<sup>3</sup>, Stephen W. Hager<sup>4</sup>,  
Andrew J. Arnsberg<sup>5</sup>, Fred Murphy<sup>6</sup>

---

U.S. GEOLOGICAL SURVEY

Open File Report 01-89

<sup>1</sup> [btopping@usgs.gov](mailto:btopping@usgs.gov), U.S. Geological Survey, Menlo Park, CA

<sup>2</sup> [kuwabara@usgs.gov](mailto:kuwabara@usgs.gov), U.S. Geological Survey, Menlo Park, CA

<sup>3</sup> [parchaso@usgs.gov](mailto:parchaso@usgs.gov), U.S. Geological Survey, Menlo Park, CA

<sup>4</sup> [swhager@usgs.gov](mailto:swhager@usgs.gov), U.S. Geological Survey, Menlo Park, CA

<sup>5</sup> [aarnsberg@usgs.gov](mailto:aarnsberg@usgs.gov), U.S. Geological Survey, Menlo Park, CA

<sup>6</sup> [fmurphy@usgs.gov](mailto:fmurphy@usgs.gov), U.S. Geological Survey, Menlo Park, CA

**U.S. DEPARTMENT OF THE INTERIOR**

**Gale A. Norton, Secretary**

**U.S. GEOLOGICAL SURVEY**

**Charles G. Groat, Director**

Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Geological Survey.

---

For addition information contact:

James S. Kuwabara  
U.S. Geological Survey  
345 Middlefield Road, MS 439  
Menlo Park, CA 94025

Copies of this report may be purchased from:

U.S. Geological Survey  
Branch of Information Services  
Box 25286  
Denver, CO 80225

## TABLE OF CONTENTS

|  |    |
|--|----|
| <a href="#">Executive Summary</a> -----                            | 5  |
| <a href="#">Background</a> -----                                   | 9  |
| <a href="#">Results and Discussion</a> -----                       | 11 |
| <a href="#">Study Design and Methods</a> -----                     | 15 |
| <a href="#">Bibliography</a> -----                                 | 17 |
| <a href="#">Acknowledgements</a> -----                             | 20 |
| Appendix 1: <a href="#">Comments on the Report Structure</a> ----- | 21 |
| Appendix 2: <a href="#">List of Figures</a> -----                  | 22 |
| Appendix 3: <a href="#">List of Tables</a> -----                   | 23 |

# **Benthic Flux of Dissolved Nickel into the Water Column of South San Francisco Bay**

by Brent R. Topping, James S. Kuwabara, Francis Parchaso, Stephen W. Hager,  
Andrew J. Arnsberg, Fred Murphy

## **Executive Summary**

Field and laboratory studies were conducted between April, 1998 and May, 1999 to provide the first direct measurements of the benthic flux of dissolved (0.2-micron filtered) nickel between the bottom sediment and water column at three sites in the southern component of San Francisco Bay (South Bay), California, ([Background, Fig. 1](#)). Dissolved nickel and predominant ligands (represented by dissolved organic carbon, and sulfides) were the solutes of primary interest. Benthic flux (sometimes referred to as internal recycling) represents the transport of dissolved chemical species between the water column and the underlying sediment.

Water-quality managers are often faced with requests to reconsider criteria for contaminant loads to aquatic systems. This is particularly true in San Francisco Bay where contaminants (e.g., nickel and copper) enter the estuarine water column from a wide range of sources (e.g., municipal and industrial discharge, urban and agricultural runoff, weathering processes, and internal remobilization). There have been frequent demands by managers and the general public to quantify the connections between fluxes of contaminants and the health, abundance, and distribution of biological resources (Kuwabara and others, 1999), motivating three decades of sustained progress by the USGS in better understanding San Francisco Bay. One relatively new field of study in San Francisco Bay focuses on a poorly understood, yet potentially predominant, source of contaminants to the bay: internal recycling or benthic flux of biologically reactive trace metals and ligands. This bioavailability and flux of contaminants and nutrients into the water column of San Francisco Bay is affected by chemical (i.e., oxidation-reduction reactions, complexation and repartitioning) and benthic biological processes (Flegal and others, 1991; Kuwabara and others, 1996; Grenz and others, 2000)

Results described herein integrate current project studies with information needs identified by State Water Resources Control Board and local stakeholders to provide initial determinations of dissolved nickel flux from the sediments into the water column of the South Bay. Recent metal-transport modeling studies by Wood and others (1995) clearly indicated the importance of non-point (including benthic) metal sources to South San Francisco Bay, although nickel was not included in that study. Quantifying and understanding the magnitude and variability of these fluxes are critical to the accurate assessment of contaminant sources and loads as well as to the development of process-integrated water-quality models for this estuary.

With a variety of strategies under consideration to determine Total Maximum Daily Loads (TMDL) of contaminants into the South Bay by point sources, the primary question posed in this study was, "Is the magnitude of the benthic flux of dissolved nickel into South Bay significant

relative to known surface-water inputs?” The question was motivated by a number of factors. First, serpentine formations near the South Bay provide a natural source of nickel that accumulates in bottom sediments. San Francisco Bay in general, and the South Bay in particular, exhibit elevated nickel concentrations in the bottom sediment (~90 - 110 micrograms per gram; Hornberger and others, 1999; Wellise and others, 1999) relative to average crustal concentrations (58 - 75 micrograms per gram; Verhoogen and others, 1970; Riley and Chester, 1971). One might therefore question whether some fraction of this sediment-associated nickel becomes remobilized for transport to the overlying water.

Second, elevated dissolved nickel concentrations in the South Bay have prompted a reexamination of the dominant sources so that appropriate TMDL strategies can be designed and implemented. Third, changes in oxidizing or reducing (redox) conditions and nutrient availability near the sediment-water interface (e.g., during phytoplankton blooms) can dramatically alter the mobility of metals and ligands associated with the bottom sediment as episodic sources of carbon settle out and accumulate (Thompson and others, 1981). Finally, there is a growing body of evidence from other aquatic systems that benthic flux or internal recycling of contaminants and nutrients is an important process to consider in developing appropriate water-quality models for San Francisco Bay (Berelson and others, 1982; Flegal and others, 1991; Kuwabara and others, 1999). The question is therefore posed to provide clarification of existing conceptual models for contaminant transport within the South Bay.

This report is formatted unconventionally in a pyramid-like structure to effectively serve the diverse group of people that want to access the information in various levels of detail ([Appendix 1](#)). The report enables quick transitions between the initial [summary information](#) (figuratively at the top of the pyramid) and later details of [methods](#) or [results](#) (that is, figuratively towards the base) using hyperlinks to supporting figures and tables, and an electronically linked [Table of Contents](#).

Sampling was performed at three U.S. Geological Survey (USGS) designated stations ([Fig. 1](#)) in the South Bay: numbers 25 (a western shoal station with fine silt/clay-textured bottom sediment), 29A (a main channel station with sediment texture similar to station 25), and 46 (an eastern shoal station with sandy bottom-sediment texture). The stations are located 10.3, 1.4 and 2.1 km south of the San Mateo Bridge, respectively (all stations are north of the Dumbarton Bridge). Samples were collected near slack tide before ebb. Therefore, the depths of the shoal stations (25 and 46) were <2 m, and the main-channel station (29A) depth varied between 13 and 14 m. Three replicate sediment cores were collected from two stations for incubation experiments to provide flux estimates and benthic biological characterizations on April 8, 1998, September 16, 1998, and May 26, 1999 ([Fig. 2](#)). Ancillary data, including nutrient and ligand fluxes, were determined to provide a water-quality framework from which to compare the dissolved nickel results. The following major observations from interdependent physical, biological, and chemical data were made:

#### Physical and Biological Characterizations

1. The surficial sediment at Stations 25 and 29A were of generally higher porosity (range of 0.80 – 0.93;  $0.88 \pm 0.04$ ) than at Station 46 (range of 0.64 to 0.83;  $0.77 \pm 0.09$ ), although the difference was not statistically significant at the 95 percent

confidence level ([Table 1](#)). This is consistent with the greater difficulty collecting cores at Station 46 because of the sandier sediment texture than the other two sites.

2. Taxonomic analyses of the macrobenthos indicated highly variable benthic communities over both temporal and spatial scales ([Table 2](#); [Macrobenthos discussion](#)). Macroinvertebrates may enhance the solute flux across the sediment-water interface due to biologically enhance advection (i.e., bioirrigation/ bioturbation as a result of feeding and sensing behaviors).
3. Chlorophyll-*a* concentrations ranged from 0.04 to 13.7 micrograms per square centimeter with highest concentrations observed at Station 25 during the spring, 1998 sampling, and the lowest concentrations at the same site during the fall sampling for that year ([Table 3](#)).
4. Benthic bacterial concentrations were of the order of  $10^8$  -  $10^9$  cells per gram-surficial sediment ([Table 4](#)). There are limited reports of bacterial abundances in estuarine sediment, but the observed values are consistent with those found in the Saint Lawrence Estuary ( $\sim 10^9$  cells per gram; Lavigne and others, 1997). Microbial communities may be critical in establishing the redox gradients that regulate the remobilization, transformation and subsequent transport of sediment-associated contaminants like nickel. Station 46 consistently exhibited lower bacterial abundance than either Stations 25 or 29A for a given sampling date.

Chemical Characterizations – A summary of observations about measured chemical parameters is provided below. Unless otherwise specified, the concentrations discussed in this section operationally refer to dissolved (i.e., 0.2 micron-filtered samples) concentrations.

1. Water-column concentrations – Dissolved macronutrient concentrations in the bottom water (approximately 1 meter above the sediment-water interface) were consistently higher (frequently by orders of magnitude) than surface-water determinations reported for similar times and locations (Regional Monitoring Program, 2001). This is consistent with measured positive benthic fluxes for the macronutrients ([Table 5](#)).

Dissolved-Ni concentrations in the bottom water over the three sampling dates ranged from a minimum of 34 (Sta. 25, Sept. 1998) to a maximum of 43 nM (Sta. 29A, May 1999). No chemical analyses for surface waters on these dates, at these sites, were available. However, when compared to Regional Monitoring Program data at similar times and locations (Regional Monitoring Program, 2001), higher dissolved-nickel concentrations at depth were only evident during two of the three sampling periods. Dissolved-nutrient data for bottom waters was only available for the spring, 1999 experiment, and indicated bottom-water concentrations that were orders of magnitude higher than surface-water concentrations during that period.

2. Benthic flux of nutrients (nitrogen species, orthophosphate) – Similar to dissolved nickel results, benthic flux of macronutrients determined from core incubations were

consistently significant (that is of equivalent or greater order of magnitude) relative to surface-water inputs ([Table 6](#), [Fig. 3](#)).

3. Benthic flux of nickel – Flux estimates from core-incubations, when areally averaged over the South Bay (554 square kilometers; Cheng and Gartner, 1985), were significant in magnitude relative to freshwater point sources (Davis and others, 1991; San Jose/Santa Clara Water Pollution Control Plant, 1999; Davis and others, 2000) ([Table 7](#), [Fig. 4](#); [Nickel discussion](#)). This is consistent with previous determinations for other metals (Flegal and others, 1991; Flegal and others 1996; Kuwabara et al, 1996), and with the potential remobilization of sediment-associated metals that have been ubiquitously distributed in the South Bay (Wellise and others, 1999).

Since the current study indicates that the magnitude of measured benthic fluxes for nickel were significant relative to major fresh water inputs, metal remobilization from the sediment should be considered in establishing future TMDL strategies for the watershed.

## **BACKGROUND** - What is benthic flux and why should it be considered?

Many fundamental processes affect the transport of dissolved chemical species (e.g., nutrients, metals, or ligands) through and within an estuary. A conceptual model of these processes ([Fig. 5](#)) illustrates some physically based processes that have been examined and carefully quantified for several years (e.g., advective transport and point source inputs) (Fisher and others, 1979).

Conversely, there are flux terms in the conceptual model that have received little attention. A prime example is the benthic flux term where no direct measurements for dissolved nickel have been available. Benthic flux (sometimes referred to as internal recycling) represents the transport of dissolved chemical species between the water column and the underlying sediment. Flux of solutes can be either positive (from the sediment into the water column) or negative (from the water column into the sediment). It can vary over multiple temporal and spatial scales.

As a result of physical, chemical, and biological changes in the vertical cross section of the sediment-water interface, geochemical gradients take on a variety of forms that have been previously reported (Kuwabara and others, 2000; [Fig. 6](#)). When the solute concentration above and below the sediment-water interface are equivalent, there is no discernable gradient, and consequently no net transfer of that substance across the interface; **no benthic flux**. In contrast, when solute concentrations in the water column are higher than those in the bottom sediment pore waters, a **negative** benthic flux results whereby the substance moves **into** the sediment. Dissolved oxygen is a typical example of such a solute where microbial respiration can create a sediment demand for oxygen. When concentrations in the water column are lower than those in sediment pore waters, the vertical concentration gradient can physically drive the **release** of dissolved chemical species from the sediment to the overlying water; a positive benthic flux. The remobilization or chemical transformation of sediment-associated trace metals or ligands may represent such gradients.

The examples above are highly simplified. When interdependent factors regulate the benthic flux of biologically reactive substances, the vertical gradient for one dissolved species may be dependent on the gradient of another chemical species. For example, an **attenuated release** may occur when solute concentrations increase below the sediment-water interface only when another solute is depleted. Dissolved iron often behaves in this manner when suboxic conditions reduce it from ferric to ferrous forms, increasing its solubility. Additionally, macroinvertebrates can biologically enhance the benthic flux by irrigating, or perturbing surficial sediment layers (**bioirrigation**, **bioturbation**, or biologically enhanced **advection**). Certain productive benthic communities can enhance benthic flux beyond diffusive-control estimates by orders of magnitude (Kuwabara and others, 1999). In summary, vertical chemical gradients generated by a variety of interdependent biogeochemical processes induce the movement of a dissolved chemical species like nickel across the sediment-water interface.

Scientists and water-quality managers are only beginning to appreciate the importance of benthic flux in many aquatic environments. In recent years (that is within the past decade or two), researchers have gradually realized that there are non-hydrologic processes (e.g., benthic flux) that must be incorporated into water-quality models to generate physically meaningful modeling results. The USGS has been involved in studies of San Francisco Bay for

approximately three decades, but as with many other disciplines, benthic-flux studies only developed within the past decade (Flegal and others, 1991; Kuwabara and others, 1996). One possible explanation for the lag in these studies relative to other transport processes is the fact that quantifying benthic flux is instrument and manpower intensive. Each flux estimate requires a concentration time series or vertical profile.

As described in the Executive Summary, the primary question posed in this study was, “Is the magnitude of the benthic flux of dissolved nickel significant relative to surface-water inputs?” Although the technical rationale for posing that question is multifaceted ([introductory comments](#)), a major objective of the study was to offer some help and scientific basis for directing TMDL strategies for the watershed by providing the first comparisons of direct benthic-flux measurements to surface water inputs.

## RESULTS AND DISCUSSION

Physical Data – There was a visible difference in sediment texture between eastern-shallow Station 46 (sandy; difficult to retain sediment cores), and both the western-shallow Station 25 and the main-channel Station 29A (noticeably higher fraction of silt and clay that facilitated core-tube penetration, and retention of sediment cores). Measured porosities were consistent with that observation and Stations 25 and 29A had higher porosities (0.80 – 0.93) than Station 46 (0.64 to 0.83; [Table 1](#)).

### Biological Data

1. Chlorophyll-*a* – Please refer directly to table summaries ([Table 3](#)).
2. Benthic bacteria – The benthic microbial community establishes redox gradients and thereby facilitates changes in the chemical structure, mobility, and biological availability of contaminants. The highest bacterial abundance was observed at Station 25 during April 1998 and was coincident with the highest sulfide fluxes for the study. This suggests an increase in sulfate-reducing bacteria with a source of organic carbon near the sediment-water interface, possibly as a consequence of the spring algal bloom. A similar range of benthic bacterial abundance (of the order of  $10^9$  cells per gram) was reported for the trough of the Saint Lawrence Estuary (Lavigne and others, 1997).
3. Benthic macroinvertebrates - At Station 46, the eastern shallow station, the dominant taxa in the benthic community changed from annelids in the spring of 1998 to bivalves in the spring of 1999 ([Fig. 7a](#)). The number of taxa, as well as the number of individuals, in the benthic community was greater in 1999 than in 1998. Combined wet weight data also reflected these trends. Annelids represented 89% of total macroinvertebrate biomass in April 1998. In contrast, mollusks represented 69% of total macroinvertebrate biomass in May 1999. However, total biomass was about five times greater in 1999 than in 1998, and annelid biomass in 1999 was greater than 1998.

At Station 25, the western shallow station, the abundance of individuals and the number of taxa were greater in September 1998 than April 1998 ([Fig 7b](#)). In April 1998, many of the common species found in South San Francisco Bay were present. In September 1998 the community was dominated by the filter feeders *Corophium* sp. and *Ampelisca abdita*. Biomass data reflected the equal contribution of crustaceans and annelids to the community in April 1998 and the increase in *Corophium* and *Ampelisca* in September 1998. *Corophium* and *Ampelisca* have been shown to enhance solute benthic flux by their feeding and sensing behaviors (Miller, 1984; Word, 1980). The increase in bivalve biomass between September and April was attributed to the increase in abundance of *Tapes japonica* in September.

Cores from Station 29A, the main channel station, were only obtained in May 1999, and the community composition and biomass was dominated by *Ampelisca abdita* (Fig. 7c).

Chemical Data – For consistency with previous geochemical studies, flux estimates are provided in molar units.

1. Bottom-water nickel concentrations – In two instances at Station 25 (spring and summer, 1998), the bottom-water concentrations (34 and 43 nanomoles per liter) indicated slightly higher concentrations relative to surface-water concentrations that were determined as part of the Regional Monitoring Program (30 and 35 nanomoles per liter, respectively) (Table 5). In contrast, the bottom-water concentration at Station 29A in May, 1999 (28 nanomoles per liter) was lower than surface-water determinations at spatially comparable sites (32 and 102 nanomoles per liter at San Bruno Shoals and Redwood Creek, respectively) in April, 1999, the closest RMP sampling date to our experiment.
2. Nickel benthic flux - The magnitude of the benthic flux for dissolved nickel displayed both small scale (within site) and large scale (between site) variability, but were consistently significant, when areally averaged (42 +/- 16 kilograms per day; Table 7), relative to municipal inputs (3 kilograms per day; San Jose/Santa Clara Water Pollution Control Plant, 1999). Previously, nickel loading to the South Bay from municipal and industrial effluents was estimated at 16 kilograms per day (Davis and others, 1991). Recent estimates by Davis and others (2000) indicate that total nickel loading to San Francisco Bay from external sources is dominated by stormwater runoff. The authors included five hydrologic units (areas) that flow into the South Bay: Alameda Creek, Fremont Bayside, Coyote Creek, Guadalupe River and Palo Alto. Two other hydrologic units, East Bay Cities and San Mateo Bayside, may partially contribute to South Bay inputs. The nickel loading associated with stormwater runoff from all seven units is approximately 56 kilograms per day.

Given ongoing nickel and copper toxicological studies in the South Bay and other parts of the estuary, it may be useful to note that the average dissolved-copper fluxes for stations 25, 29A, and 46 were 7, -1 and -23 nanomoles per square meter per hour, considerably lower than observed nickel fluxes, and distinct because of variable direction.

3. Nutrients - The benthic flux of macronutrients determined in these experiments are comparable to measurements made in 1996, (Grenz and others, 2000), but are considerably lower than measurements made between 1991 and 1993 (Caffrey and others, 1996). This trend is consistent with long-term reductions in phosphorus loading to the South Bay (Schemel and others, 1999). Although the results of this study only provide initial estimates of spatial and temporal variability of these fluxes, measurements reported in this and previous studies suggest that internal recycling of

nutrients in the South Bay needs to be considered when developing models that affect water-quality management decisions for the South Bay.

Because orthophosphate has a high affinity to adsorb onto metal-oxide surfaces under oxic, pH-neutral conditions (Kuwabara, 1992), molar nitrogen to phosphorus (N to P) flux ratios for the South Bay may have particular significance. The molar N to P flux ratios for Stations 29A and 46 in May, 1999, the only date among the three when nutrient concentrations are available, are 19 and 26, respectively (Fig. 8). By comparison, the molar N to P Redfield Ratio is only 16 (7 by mass ratio; Redfield and others, 1963). That is, on average, approximately 16 moles of nitrogen (220 grams) are taken up for every mole of phosphorus (31 grams) to generate tissue of primary producers. Consistent with other aquatic systems (Kuwabara and others, 2000), the N to P ratios determined for both sites suggest that the sediments can promote the recycling of nitrogen while retarding sediment sources of biogenically derived phosphorus into the water column by sorption/desorption reactions. In contrast, the average molar N to P loading ratio in 1998 for the San Jose/Santa Clara Water Pollution Control Plant was approximately 5 (San Jose/Santa Clara Water Pollution Control Plant, 1999), considerably below the Redfield Ratio. This suggests that surface-water inputs may act as a source of phosphorus for primary production in the South Bay.

Time series of dissolved nutrients from incubated cores indicate that concentration increases in the overlying water primarily occurred in the initial hours of the incubation. The strong initial flux followed by diminished changes in the overlying water chemistry may be due to a shift in the pore-water/bottom-water concentration gradient during the incubation (i.e., that a chemical equilibrium between the pore water and overlying water in certain cores was reached rapidly relative to the entire period of incubation.) This phenomenon occurred with more frequency at Station 46 and is consistent with the lower bottom-water concentrations there relative to the other two sites. Such nonlinearities generated lower flux estimates and coefficients of determination when data from the entire incubation were used in the flux determination as opposed to just to the initial hours (Table 8). However, even if data over the entire incubation are used in the flux calculation, the areally averaged benthic flux of orthophosphate and nitrogen species are still significant relative to municipal point source inputs (Table 6). For example, the areally averaged orthophosphate fluxes from Stations 29A and 46 cores were 1,440 and 2,470 kilograms per day, respectively, compared with a freshwater loading of approximately 1,960 kilograms per day, primarily from a major freshwater point source (San Jose/Santa Clara Water Pollution Control Plant, 1999; Table 6, Fig. 3). Ammonia benthic flux averaged approximately 6,460 and 13,330 kilograms per day from Stations 29A and 46, respectively, compared to only 490 kilograms per day from the same point source.

4. Dissolved organic carbon (DOC) - DOC concentrations serve as a model for the abundance of organic ligands that can complex and enhance the solubility of metals. Certain dissolved metals, in particular nickel, predominantly exist as organic complexes in many aquatic systems (Mantoura and others, 1978; McKnight and

others, 1983, Donat and others, 1994). The benthic flux of DOC exhibited both small-scale (within-site) and large-scale (between-site) variability, ranging from – 239 to 676 micromoles per square meter per hour ([Table 9](#)).

Bottom-water DOC concentrations were highest in summer 1998 (St. 25, 341 micromoles per liter) and lowest in spring 1999 (St. 29A, 141 micromoles per liter). When compared with RMP surface-water data, bottom-water DOC concentrations showed a vertical gradient (higher concentration at depth) in only two of three sites, similar to nickel. In contrast to many of the dissolved inorganic substances, DOC benthic fluxes did not exhibit consistent trends.

5. Dissolved Sulfides – Dissolved sulfides represent a metastable ligand with high affinity to complex most divalent metals like nickel (Hogfeldt, 1983). Sulfide flux was consistently positive (that is, from the sediment to the overlying water column) and ranged from 181 to 752 nanomoles per square meter per day. The highest sulfide fluxes were observed in April 1998, and were significantly higher than the other two dates ([Table 10](#)). Notably, the highest bacterial abundances also were observed during that sampling event.

## STUDY DESIGN AND METHODS

The protocol described in this section focuses on method applications in this series of three core-incubation experiments. Details (e.g., quality control specifications) for each analysis has been previously documented (Woods and others, 1999).

### Coring Operation

On each of three sampling days (4/8/98, 9/16/98, 5/26/99), four cores were collected at each of two sites using a coring device fabricated from non-metallic parts (Savillex Corporation, Minnetonka, Minnesota). To avoid sample contamination, wetted surfaces of the coring device and core tubes were acid-washed polypropylene or fluoroethylene polymer. The eight cores were aerated overnight to condition wetted surfaces before the incubation period ([Fig. 2](#), [Fig. 9](#)). Based on visual determination of any disturbances of the sediment-water interface, three of the four cores per site were selected for incubation. No cores were taken 9/16/98 at station 46 due to operator error. Therefore, only four suitable cores, all from station 25, were collected and incubated for this date. Water overlying the sediments in the selected cores was sampled at four intervals during a 12-hour incubation. Trace element and nutrient samples were processed in a Class-100 laminar-flow hood. The concentration time series was used to determine the benthic flux from each of the incubated cores.

### Physical Data

Sediment Porosity – After core-incubations, approximately 10 milliliters of surficial sediment was collected from each core, then bottled, and refrigerated in darkness. Wet weight and dry weight after lyophilization was measured to calculate porosity ([Table 1](#)).

### Biological Data

1. Benthic Macrofauna – After core incubations, each core was sieved (500-micron mesh). The sieved samples were fixed with 10% buffered formalin, transferred to 70% ethanol, then sorted and identified to the lowest practicable taxonomic level ([Fig. 10](#)).
2. Benthic Chlorophyll-*a* - Each incubated core was sub-sampled in triplicate for benthic chlorophyll-*a*. Surficial sediment was collected on a glass-fiber filter and buffered with magnesium carbonate. Samples were then frozen in darkness until spectrophotometrically analyzed (Thompson and others, 1981; Franson, 1985) ([Fig. 11](#)).

3. Benthic Bacteria - After core incubations, each core was sub-sampled using a syringe corer. Approximately 10 milliliters of surficial sediment was collected and fixed with 0.5 milliliters of 37% formalin. Samples were then refrigerated in darkness. Benthic bacterial concentrations were determined by direct epifluorescent counts using acridine-orange preparations.

### Chemical Parameters

1. Bottom-water sampling – Prior to coring at each sampling site, water-column samples were collected for trace metals analyses primarily using inductively coupled plasma mass spectrometry. In May, 1999, samples were also analyzed for dissolved (0.2-micron filtered) macronutrients by automated spectrophotometry. Samples were pumped from approximately 1 meter above the bottom using a high-displacement peristaltic pump and a tethered length of rigid fluoroethylene polymer tubing.
2. Nutrients (nitrogen species and orthophosphate) – Dissolved nutrients from water-column sampling, and core incubations were filtered with 0.2 micron polycarbonate filters. The filtered samples were immediately refrigerated in darkness, but unlike trace metal samples, were not acidified. Nutrient analyses were performed by automated spectrophotometry ([Fig. 12](#)).
3. Dissolved Nickel – Nickel samples from water-column sampling, and core incubations were filtered with 0.2 micron acid-washed polycarbonate filters. Filtered samples were acidified with quartz-distilled nitric acid to 0.03N, and refrigerated in darkness until analyses. Metal analyses were performed by flow-injection inductively coupled plasma mass spectrometry (FIAS ICP-MS) using external standardization ([Fig. 13](#)).
4. Dissolved organic carbon (DOC) – Dissolved organic carbon was determined by high-temperature non-catalytic combustion (Qian and Mopper, 1997). Potassium phthalate was used as the standard. Low-DOC water (blanks <40 micrograms-organic C per liter) was generated from a double-deionization unit with additional ultraviolet treatment (Milli-Q Gradient – Millipore Corporation) ([Fig. 14](#)).
5. Sulfides – Dissolved sulfides in overlying-water samples were analyzed by square-wave voltammetry (Kuwabara and Luther, 1993) ([Fig. 15](#)).

## REFERENCES CITED

- Berelson, W. M., Hammond, D. E., and Fuller, C., 1982, Radon-222 as a tracer for mixing in the water column and benthic exchange in the southern California borderland: *Earth and Planetary Science Letters*, v. 61, p. 41-54.
- Caffrey, J., Hammond, D., Kuwabara, J., Miller, L., and Twilley, R., 1996, Benthic processes in San Francisco Bay: the role of organic inputs and bioturbation, in Hollibaugh, J.T., ed., *San Francisco Bay: the Ecosystem: American Association for the Advancement of Science, Pacific Division, San Francisco*, p. 425-442.
- Cheng, R.T. and Gartner, J.W., 1985, Harmonic analysis of tides and tidal current in South San Francisco Bay, California: *Estuarine, Coastal and Shelf Science*, v. 21, p. 57-74.
- Davis, J.A., Gunther, A.J., Richardson, B.J., O'Connor, J.M., Spies, R.B., Wyatt, E., Larson, E., and Meiorin, E.C., 1991, Status and trends report on pollutants in the San Francisco Estuary: San Francisco Estuary Project, Richmond, CA, 240 pp.
- Davis, J.A., McKee, L.J., Leatherbarrow, J.E., and Daum, T.H., 2000, Contaminant loads from stormwater to coastal waters in the San Francisco Bay Region, San Francisco Estuary Institute, Richmond, CA, 82 pp. <http://www.sfei.org/rmp/reports/AB1429/ab14.html>
- Donat, J.R., Lao, K.A., and Bruland, K.W., 1994, Speciation of dissolved copper and nickel in South San Francisco Bay: a multi-method approach. *Analytica Chimica Acta* 284: 547-571.
- Fischer, H.B., List, E.J., Koh, R.C.Y., Imberger, J., and Brooks, N.H., 1979, *Mixing in Inland and Coastal Waters: Academic Press, Inc., Orlando, FL*, 483 p.
- Flegal, A.R., Smith G.J., Gill, G.A., Sanudo-Wilhelmy, S., Scelfo, G., and Anderson, L.C.D., 1991, Dissolved trace element cycles in the San Francisco Bay estuary: *Marine Chemistry*, v. 36, p. 329-363.
- Flegal, A.R., Rivera-Duarte, I., Ritson, P.I., Scelfo, G.M., Smith, G.J., Gordon, M.R., and Sanudo-Wilhelmy, S.A., 1996, Metal contamination in San Francisco Bay waters: Historic perturbations, contemporary concentrations, and future considerations, in Hollibaugh, J.T., ed., *San Francisco Bay: the Ecosystem: American Association for the Advancement of Science, Pacific Division, San Francisco*, p. 173 – 188.
- Franson, M.A.H., 1995, *Standard Methods for the Examination of Water and Wastewater, Sixteenth Edition, Method 1003C.6: American Public Health Association, American Water Works Association, Water Pollution Control Federation, Washington, D.C.*, 1268 p.
- Grenz, C., Cloern, J.E., Hager, S.W., and Cole, B.E., 2000, Dynamics of nutrient cycling and related benthic nutrient and oxygen fluxes during a spring phytoplankton bloom in South San Francisco Bay (USA): *Marine Ecological Progress Series*, v. 197, p. 67-80.

- Hogfeldt, E., 1983, Stability constants of Metal-Ion Complexes. Part A, Inorganic Ligands. Pergamon Press, Oxford. 310 pp.
- Hornberger, M.I., Luoma, S.N., van Geen, A., Fuller, C., Anima, R., 1999, Historical trends of metals in the sediments of San Francisco Bay, California: *Marine Chemistry*, v. 64, p. 39-55.
- Kuwabara, J.S., 1992, Associations between benthic flora and diel changes in dissolved arsenic, phosphorus, and related physio-chemical parameters: *Journal of the North American Benthological Society*, v. 11, p. 218-228.
- Kuwabara, J.S. and Luther, G.W., III, 1993, Dissolved sulfides in the oxic water column of San Francisco Bay, California: *Estuaries*, v. 16, p. 567-573.
- Kuwabara, J.S., Chang, C.C.Y., Khechfe, A.I. and Hunter, Y.R., 1996, Implications of dissolved sulfides and organic substances for the chemical speciation of trace contaminants in the water column of San Francisco Bay, California, in Hollibaugh, J.T., ed., *San Francisco Bay: the Ecosystem: American Association for the Advancement of Science, Pacific Division, San Francisco*, p. 157-172.
- Kuwabara, J.S., van Geen, A., McCorkle, D.C., and Bernhard, J.M., 1999, Dissolved sulfide distributions in the water column and sediment pore waters of Santa Barbara Basin: *Geochimica et Cosmochimica Acta*, v. 63, no. 15, p. 2199-2209.
- Kuwabara, J.S., Berelson, W.M., Balistrieri, L.S., Woods, P.F., Topping, B.R., Steding, D. J., and Krabbenhoft, D.P., 2000, Benthic flux of metals and nutrients into the water column of lake Coeur d'Alene, Idaho: Report of an August, 1999 Pilot Study: USGS Water-Resources Investigations Report 00-4132, 74 p. (Internet access at: <http://water.usgs.gov/pubs/wri/wri004132>)
- Lavigne, C., Juniper, S.K., and Silverberg, N., 1997, Spatio-temporal variability in benthic microbial activity and particle flux in the Laurentian Trough: *Deep-Sea Research I*, v. 44, p. 1793-1813.
- Mantoura, R.F.C., Dixon, A., and Riley, J.P., 1978, The complexation of metals with humic materials in natural waters: *Estuarine and Coastal Marine Sciences*, v. 6, p. 387-408.
- McKnight, D.M., Feder, G.L., Thurman, E.M., Wershaw, R.L. and Westall, J.C., 1983, Complexation of copper by aquatic humic substances from different environments: *Science of the Total Environment*, v. 28, p. 65-76.
- Miller, D.C., 1984, Mechanical post-capture particle selection by suspension- and deposit-feeding Corophium: *Journal of Experimental Biology and Ecology*, v. 82, p. 59-76.
- Qian, J. and Mopper, K., 1996, Automated high-performance, high-temperature combustion total organic carbon analyzer: *Analytical Chemistry*, v. 68, p. 3090-3097.

- Redfield, A.C., Ketchum, B.H., and Richards, F.A., 1963, The influence of organisms on the composition of sea-water, in Hill, M.N., ed., *The Sea*: Wiley-Interscience, New York, v. 2, p. 26-77.
- Regional Monitoring Program (RMP), 2001, website: <http://www.sfei.org/rmp/rmpwater.htm>
- Riley, J.P. and Chester, R., 1971, *Introduction to Marine Chemistry*: Academic Press, London, England, 465 p.
- San Jose/Santa Clara Water Pollution Control Plant, 1999, Combined December 1998 and Annual 1998 Self-Monitoring Report: City of San Jose, California, 34 p.
- Schemel, L.E., Hager, S.W., and Peterson, D.H., 1999, Reduced phosphate loading to South San Francisco Bay, California: Detection of effects in the water column, In Morganwalp, D.W., and Buxton, H.T., eds., *U.S. Geological Survey Toxic Substances Hydrology Program--Proceedings of the Technical Meeting, Charleston, South Carolina, March 8-12, 1999--Volume 2--Contamination of Hydrologic Systems and Related Ecosystems: U.S. Geological Survey Water-Resources Investigations Report 99-4018B*, p. 59-64.
- Thompson, J.K., Nichols, F.H., and Wienke, S.M., 1981, Distribution of benthic chlorophyll in San Francisco Bay, California, February 1980 – February 1981: U.S. Geological Survey Open File Report 81-1134, 55 p.
- Verhoogen, J., Turner, J., Weiss, L.E., Wahrhaftig, C., and Fyfe, W.S., 1970, *The Earth: An Introduction to Physical Geology*: Holt, Rinehart and Winston, Inc., Austin, Texas, 748 p.
- Wellise, C.J., Luoma, S.N., Cain, D./J., Brown, C., Hornberger, M., Bouse, R., 1999, Near field receiving water monitoring of trace metals in clams (*Macoma balthica*) and sediments near the Palo Alto and San Jose/Sunnyvale Water Quality Control Plants in South San Francisco Bay, California: 1998: U.S. Geological Survey Open File Report 99-455, 101 p.
- Wood, T.M., Baptista, A.M., Kuwabara, J.S. and Flegal, A.R., 1995, Diagnostic modeling of trace metal partitioning in south San Francisco Bay: *Limnology and Oceanography*, v. 40, p. 345-358.
- Woods, P.F., Nearman, M.J., and Barton, G.J., 1999, Quality Assurance Project Plan for U.S. Geological Survey Studies in Support of Spokane River Basin RI/FS.: U.S. Environmental Protection Agency, Seattle, Washington, and U.S. Geological Survey, Boise, Idaho, 153 p.
- Word, J.Q., 1980, Classification of benthic invertebrates into infaunal trophic index feeding groups, Bascom, W., ed., *Biennial Report: Southern California Coastal Research Project*, Long Beach, CA, p. 103-121.

## **Acknowledgements**

The authors are grateful to J. Carter and S. Fend for critical analyses of benthic-invertebrate samples. Logistical and facilities support by C. Azevedo and B. Richards was also much appreciated. Discussions and advice from J. Thompson, and reviews by M. Hornberger and J. Carter were critical to the development of this report. The California State Water Resources Control Board and P. Mulvey are gratefully acknowledged for their support of this work, as is the U.S. Geological Survey Toxic Substances Hydrology Program for long-term support of this area of research.

Product names are provided for identification purposes only and do not constitute endorsement by the U.S. Geological Survey.

## **Appendix 1: Comments on the Report Structure**

A major objective of this electronic document is to provide a structure that is easily accessible to a potentially wide range of interests in this work. Agency reports in the form of electronically linked products are a fairly recent alternative, but pathways within this document have been constructed to be both logical and intuitive. In addition to hyperlinks within the document to supporting figures and tables, Appendices 2 and 3 provide a quick way to directly review and examine all figures and tables.

Although hard copies of this report are available on request, the advantages of the electronic version relative to the hard copy are substantial in many respects, but particularly in speed of information access at multiple levels of detail.

Your comments about how to improve this evolving type of product are most welcome and may be directed to any of the email addresses provided in the title page, but we request that at least a copy of all comments about the report be addressed to the major author ([btopping@usgs.gov](mailto:btopping@usgs.gov)) so that they may be compiled for future revisions and reports.

## Appendix 2: List of Figures

- [Fig. 1](#) – Sampling/Coring Sites in the South Bay
- [Fig. 2](#) – Coring Operation
- [Fig. 3](#) – Dissolved-nutrients Flux Comparison
- [Fig. 4](#) – Dissolved-nickel Flux Comparison
- [Fig. 5](#) – Conceptual Model of Solute Transport
- [Fig. 6](#) – Processes Regulating Benthic Flux
- [Fig. 7a](#) – Macrobenthos counts at eastern shallow station (46)
- [Fig. 7b](#) – Macrobenthos counts at western shallow station (25)
- [Fig. 7c](#) – Macrobenthos counts at main channel station (29A)
- [Fig. 8](#) – Dissolved Nitrogen to Phosphorous (N:P) Benthic Flux Ratio
- [Fig. 9](#) – Incubation Core Design
- [Fig. 10](#) – Macroinvertebrate Taxonomy
- [Fig. 11](#) – Benthic Chlorophyll Analyses
- [Fig. 12](#) – Spectrophotometric Nutrient Analyses
- [Fig. 13](#) – Dissolved-nickel Analyses by ICP-MS
- [Fig. 14](#) – Dissolved Organic Carbon (DOC) Analyses
- [Fig. 15](#) – Dissolved sulfides by Square-wave Voltammetry

### Appendix 3: List of Tables

|                          |  |
|--------------------------|--|
| <a href="#">Table 1</a>  | – Surficial-sediment Porosities                              |
| <a href="#">Table 2</a>  | – Benthic-biological Indices                                 |
| <a href="#">Table 3</a>  | – Benthic Chlorophyll  |
| <a href="#">Table 4</a>  | – Surficial Benthic-bacteria Concentrations                  |
| <a href="#">Table 5</a>  | – Dissolved Chemical Species in the Water Column             |
| <a href="#">Table 6</a>  | – Nutrient flux from Core Incubations                        |
| <a href="#">Table 7</a>  | – Nickel flux from Core Incubations                          |
| <a href="#">Table 8</a>  | – Flux calculations using only first two hours of incubation |
| <a href="#">Table 9</a>  | – Dissolved Organic Carbon flux from Core Incubations        |
| <a href="#">Table 10</a> | – Sulfide flux from Core Incubations                         |

Fig. 1

# Benthic-flux Coring Sites




Fig. 2

## Coring Operation



Release



Retrieval



Removal

Fig. 3

# Nutrient Comparison




Fig. 4

# Nickel Comparison




Fig. 5

## Conceptual Model of Solute Transport




Fig. 6

## Processes Regulating Benthic Flux




Fig. 7a

# Macroinvertebrate Densities at Eastern Shallow Station 46

Note: If there is no bar, the corresponding species was not found in those samples.




Fig. 7b

# Macroinvertebrate Densities at Western Shallow Station 25

Note: If there is no bar, the corresponding species was not found in those samples.




Fig. 7c

# Macroinvertebrate Densities at Main Channel Station 29A

Note: If there is no bar, the corresponding species was not found in those samples.




Fig. 8

# Nitrogen to Phosphorus Ratio




Fig. 9

## Incubation Core Design




Fig. 10

## Benthic Macroinvertebrate Taxonomy



Fig. 11

## Benthic Chlorophyll Analyses



Fig. 12

## Spectrophotometric Nutrient Analyses



Fig. 13

## Dissolved-nickel Analyses by ICP-MS



Fig. 14

## Dissolved Organic Carbon (DOC) Analyses by High-Temperature Non-Catalytic Oxidation



Fig. 15



**Table 1.** Surficial-sediment Porosities

| <b>4/8/98</b> |      |             |  |                   |
|---------------|------|-------------|--|-------------------|
| Station       | Core | Porosity    | Bulk Density<br>(g-dry/cm <sup>3</sup> -sed) | Wet/dry<br>weight |
| 25            | 6    | <b>0.85</b> | 0.33   | 3.68              |
| 25            | 7    | <b>0.92</b> | 0.17   | 6.56              |
| 25            | 8    | <b>0.95</b> | 0.12   | 9.38              |
| 46            | 1    | <b>0.83</b> | 0.38   | 3.28              |
| 46            | 3    | <b>0.81</b> | 0.43   | 2.93              |
| 46            | 4    | <b>0.83</b> | 0.41   | 3.11              |

| <b>9/16/98</b> |      |             |  |                   |
|----------------|------|-------------|--|-------------------|
| Station        | Core | Porosity    | Bulk Density<br>(g-dry/cm <sup>3</sup> -sed) | Wet/dry<br>weight |
| 25             | 1    | <b>0.93</b> | 0.15   | 7.45              |
| 25             | 2    | <b>0.88</b> | 0.27   | 4.39              |
| 25             | 3    | <b>0.91</b> | 0.20   | 5.66              |
| 25             | 4    | <b>0.87</b> | 0.26   | 4.48              |

| <b>5/26/99</b> |      |             |  |                   |
|----------------|------|-------------|--|-------------------|
| Station        | Core | Porosity    | Bulk Density<br>(g-dry/cm <sup>3</sup> -sed) | Wet/dry<br>weight |
| 29A            | 2    | <b>0.80</b> | 0.47   | 4.41              |
| 29A            | 3    | <b>0.85</b> | 0.37   | 5.39              |
| 29A            | 4    | <b>0.83</b> | 0.43   | 4.79              |
| 46             | 6    | <b>0.79</b> | 0.48   | 4.37              |
| 46             | 7    | <b>0.73</b> | 0.61   | 3.55              |
| 46             | 8    | <b>0.64</b> | 0.82   | 2.97              |

**Table 2. Macrobenthos**

Mean number of individuals per square meter based on core surface area of 77 square centimeters.

| Note: Species are listed below in the same order as on the plots | Click on station name to see corresponding plot |           |             |           |           |
|--|---|-----------|-------------|-----------|-----------|
|  | Station: 46                                     |           | Station: 25 |           | Sta: 29A  |
|  | 08-Apr-98                                       | 25-May-99 | 08-Apr-98   | 15-Sep-98 | 25-May-99 |
| Phylum Mollusca  |   |           |             |           |           |
| Class Bivalvia   |   |           |             |           |           |
| <i>Potamocorbula amurensis</i>                                   | 520   |           | 303         | 195       |           |
| <i>Musculista senhousia</i>                                      |   |           | 43          | 130       |           |
| <i>Mya arenaria</i>  | 43  | 1342      | 43          | 32        |           |
| <i>Theora fragilis</i>   |   |           |             | 32        |           |
| <i>Gemma gemma</i>   | 43  | 2035      |             |           |           |
| <i>Macoma balthica</i>   | 87  |           |             |           |           |
| <i>Tapes japonica</i>  |   | 43        | 87          | 714       |           |
| Class Gastropoda   |   |           |             |           |           |
| <i>Odostomia</i> sp.   |   | 43        |             |           |           |
| Phylum Arthropoda  |   |           |             |           |           |
| Class Crustacea  |   |           |             |           |           |
| Order Amphipoda  |   |           |             |           |           |
| Family Gammaridae  |   |           |             |           |           |
| <i>Grandidierella japonica</i>                                   | 87  |           |             |           |           |
| Family Ampeliscidae  |   |           |             |           |           |
| <i>Ampelisca abdita</i>  | 87  | 87        | 1212        | 3150      | 4027      |
| Family Corophidae  |   |           |             |           |           |
| <i>Corophium acherusicum</i>                                     |   | 43        |             |           | 130       |
| <i>Corophium heteroceratum</i>                                   |   |           |             |           | 43        |
| <i>Corophium</i> sp.   |   | 43        |             | 5813      |           |
| Order Cumacea  |   |           |             |           |           |
| <i>Cumella vulgaris</i>  | 390   | 87        | 953         | 32        |           |
| <i>Nippoleucon hinumensis</i>                                    |   |           |             |           | 87        |
| Order Isopoda  |   |           |             |           |           |
| <i>Synidotea</i> spp.  | 43  |           |             |           | 87        |
| <i>Eusarsiella zostericola</i>                                   |   | 303       |             |           | 87        |
| Subclass Ostracoda   |   |           |             |           |           |
| <i>Cylindroleberis</i> sp.                                       | 43  |           | 43          | 65        |           |
| Phylum Annelida  |   |           |             |           |           |
| Class Oligochaeta  |   |           |             |           |           |
| Family Tubificidae sp.   |   | 43        |             |           | 909       |
| Class Polychaeta   |   |           |             |           |           |
| Family Capitellidae  |   |           |             |           |           |
| Capitellidae *   | 0   |           | 0           | 0         |           |
| <i>Heteromastus filiformis</i>                                   |   | 563       |             |           |           |
| Family Eunicidae   |   |           |             |           |           |
| Eunicidae *  |   |           |             | 0         |           |
| <i>Marphysa</i> nr. <i>Sanguinea</i>                             |   | 43        |             |           |           |
| Family Goniadidae  |   |           |             |           |           |
| <i>Glycinde polygnatha</i>                                       |   | 130       |             |           |           |
| <i>Glycinde</i> sp.  |   |           |             |           | 43        |
| Family Maldanidae  |   |           |             |           |           |
| Maldanidae *   | 0   |           | 0           | 43        |           |
| <i>Sabaco elongatus</i>  |   | 87        |             |           | 87        |
| Family Nephtyidae  |   |           |             |           |           |
| <i>Nephtys caecoides</i>   |   |           |             |           | 43        |
| Family Nereidae  |   |           |             |           |           |
| Nereidae *   | 0   |           |             | 39        |           |
| <i>Neanthes succinea</i>   |   | 173       |             |           |           |
| Family Polynoidae  |   |           |             |           |           |
| <i>Harmothoe imbricata</i>                                       |   |           |             |           | 43        |
| Family Spionidae   |   |           |             |           |           |
| <i>Polydora ligni</i>  |   | 43        |             |           |           |
| <i>Streblospio benedicti</i>                                     |   | 173       |             |           |           |
| Family Syllidae  |   |           |             |           |           |
| <i>Sphaerosyllis californiensis</i>                              |   |           |             |           | 43        |
| Phylum Cnidaria  |   |           |             |           |           |
| <i>Diadumene</i> sp.   |   |           |             | 32        |           |
| Phylum Tunicata  |   |           |             |           |           |
| <i>Molgula manhattensis</i>                                      |   |           | 43          | 32        |           |

\* Within these rows, zeros represent samples where the presence of worm fragments prevented identification beyond the family level. These values are thus excluded from quantification.

| Combined wet wts. grams | Station: 46 |           | Station: 25 |           | Sta. 29A  |
|-------------------------|-------------|-----------|-------------|-----------|-----------|
|                         | 08-Apr-98   | 25-May-99 | 08-Apr-98   | 15-Sep-98 | 25-May-99 |
| Molluscs                | 0.021       | 0.737     | 0.004       | 0.672     |           |
| Crustaceans             | 0.002       | 0.002     | 0.084       | 0.021     | 0.025     |
| Annelids                | 0.186       | 0.324     | 0.049       | 0.531     | 0.003     |
| Cnidaria                |             |           |             |           |           |
| Tunicata                |             |           | 0.084       | 0.063     |           |

**Table 3.** Discrete Benthic Chlorophyll Analysis

| <b>4/8/98</b> |      | Chlorophyll-a              |                            |
|---------------|------|----------------------------|----------------------------|
| Station       | Core | Mean<br>ug/cm <sup>2</sup> | S.D.<br>ug/cm <sup>2</sup> |
| 25            | 6    | <b>2.26</b>                | 0.11                       |
| 25            | 7    | <b>10.96</b>               | 2.35                       |
| 25            | 8    | <b>6.82</b>                | 5.39                       |
| 46            | 1    | <b>2.27</b>                | 1.52                       |
| 46            | 3    | <b>3.92</b>                | 0.28                       |
| 46            | 4    | <b>2.76</b>                | 0.78                       |

| <b>9/16/98</b> |      | Chlorophyll-a              |                            |
|----------------|------|----------------------------|----------------------------|
| Station        | Core | Mean<br>ug/cm <sup>2</sup> | S.D.<br>ug/cm <sup>2</sup> |
| 25             | 1    | <b>0.27</b>                | 0.11                       |
| 25             | 2    | <b>0.60</b>                | 0.32                       |
| 25             | 3    | <b>0.04</b>                | 0.06                       |
| 25             | 4    | <b>0.62</b>                | 0.15                       |

| <b>5/26/99</b> |      | Chlorophyll-a              |                            |
|----------------|------|----------------------------|----------------------------|
| Station        | Core | Mean<br>ug/cm <sup>2</sup> | S.D.<br>ug/cm <sup>2</sup> |
| 29A            | 2    | <b>0.81</b>                | 0.34                       |
| 29A            | 3    | <b>2.55</b>                | 1.29                       |
| 29A            | 4    | <b>2.03</b>                | 1.67                       |
| 46             | 6    | <b>0.59</b>                | 0.19                       |
| 46             | 7    | <b>1.14</b>                | 0.86                       |
| 46             | 8    | <b>0.56</b>                | 0.11                       |

S.D.=Standard deviation

**Table 4** Benthic Bacteria Concentrations for Core IncubationsValues in  $10^8$  bacteria cells per gram of core material

| <b>4/8/98</b> |      | Bacteria concentrations<br>$10^8$ cells/g-core |         |    |  |
|---------------|------|--|---------|----|--|
| Station       | Core | mean   | CI      | n  |  |
| 25            | 6    | <b>20.2</b>                                    | +/- 2.0 | 48 |  |
| 25            | 7    | <b>17.3</b>                                    | +/- 2.0 | 48 |  |
| 25            | 8    | <b>31.0</b>                                    | +/- 3.5 | 48 |  |
| 46            | 1    | <b>15.8</b>                                    | +/- 1.9 | 48 |  |
| 46            | 3    | <b>13.9</b>                                    | +/- 1.7 | 48 |  |
| 46            | 4    | <b>6.8</b>                                     | +/- 0.8 | 48 |  |

| <b>9/16/98</b> |      | Bacteria concentrations<br>$10^8$ cells/g-core |         |    |  |
|----------------|------|--|---------|----|--|
| Station        | Core | mean   | CI      | n  |  |
| 25             | 1    | <b>5.6</b>                                     | +/- 0.7 | 48 |  |
| 25             | 2    | <b>3.9</b>                                     | +/- 0.3 | 48 |  |
| 25             | 3    | <b>4.8</b>                                     | +/- 0.5 | 48 |  |
| 25             | 4    | <b>4.1</b>                                     | +/- 0.4 | 48 |  |

| <b>5/26/99</b> |      | Bacteria concentrations<br>$10^8$ cells/g-core |         |    |  |
|----------------|------|--|---------|----|--|
| Station        | Core | mean   | CI      | n  |  |
| 29A            | 2    | <b>12.5</b>                                    | +/- 0.8 | 48 |  |
| 29A            | 3    | <b>10.4</b>                                    | +/- 1.0 | 48 |  |
| 29A            | 4    | <b>12.8</b>                                    | +/- 0.9 | 48 |  |
| 46             | 6    | <b>5.0</b>                                     | +/- 0.6 | 48 |  |
| 46             | 7    | <b>4.1</b>                                     | +/- 0.4 | 48 |  |
| 46             | 8    | <b>3.1</b>                                     | +/- 0.4 | 48 |  |

CI=95% confidence interval

n=number of samples

**Table 5** Water Column: Comparison between bottom water and comparable surface water

| <b>Nickel</b> |         | RMP data |         | Bottom water data |      |  |
|---------------|---------|----------|---------|-------------------|------|--|
| Sta.          | Date    | Ni (nM)  | Ni (nM) | Date              | Sta. |  |
| RC            | 4/22/98 | 30       | 34      | 4/8/98            | 25   |  |
|               |         |          | 30      | 4/8/98            | 46   |  |
| RC            | 7/20/98 | 35       | 43      | 9/16/98           | 25   |  |
| RC&SB         | 4/12/99 | 67       | 28      | 5/26/99           | 29A  |  |
|               |         |          | 33      | 5/26/99           | 46   |  |

| <b>Ammonia</b> |            | RMP data   |       | Bottom water data |     |  |
|----------------|------------|------------|-------|-------------------|-----|--|
| Date           | NH4 (mg/L) | NH4 (mg/L) | Date  | Sta.              |     |  |
| RC             | 4/22/98    | 0.02       |       | 4/8/98            | 25  |  |
|                |            |            |       | 4/8/98            | 46  |  |
| RC             | 7/20/98    | 0.03       |       | 9/16/98           | 25  |  |
| RC&SB          | 4/12/99    | 0.04       | 11.14 | 5/26/99           | 29A |  |
|                |            |            | 0.93  | 5/26/99           | 46  |  |

| <b>Nitrate</b> |            | RMP data   |       | Bottom water data |     |  |
|----------------|------------|------------|-------|-------------------|-----|--|
| Date           | N03 (mg/L) | N03 (mg/L) | Date  | Sta.              |     |  |
| RC             | 1/27/98    | 0.6        |       |                   |     |  |
| RC             | 4/22/98    | n/a        |       | 4/8/98            | 25  |  |
|                |            |            |       | 4/8/98            | 46  |  |
| RC             | 7/20/98    | 0.3        |       | 9/16/98           | 25  |  |
| RC&SB          | 4/12/99    | 0.1        | 23.99 | 5/26/99           | 29A |  |
|                |            |            | 11.86 | 5/26/99           | 46  |  |

| <b>Nitrite</b> |            | RMP data   |      | Bottom water data |     |  |
|----------------|------------|------------|------|-------------------|-----|--|
| Date           | N02 (mg/L) | N02 (mg/L) | Date | Sta.              |     |  |
| RC             | 4/22/98    | 0.037      |      | 4/8/98            | 25  |  |
|                |            |            |      | 4/8/98            | 46  |  |
| RC             | 7/20/98    | 0.012      |      | 9/16/98           | 25  |  |
| RC&SB          | 4/12/99    | 0.004      | 1.18 | 5/26/99           | 29A |  |
|                |            |            | 0.18 | 5/26/99           | 46  |  |

| <b>Phosphate</b> |            | RMP data   |      | Bottom water data |     |  |
|------------------|------------|------------|------|-------------------|-----|--|
| Date             | PO4 (mg/L) | PO4 (mg/L) | Date | Sta.              |     |  |
| RC               | 4/22/98    | 0.06       |      | 4/8/98            | 25  |  |
|                  |            |            |      | 4/8/98            | 46  |  |
| RC               | 7/20/98    | 0.24       |      | 9/16/98           | 25  |  |
| RC&SB            | 4/12/99    | 0.05       | 3.86 | 5/26/99           | 29A |  |
|                  |            |            | 3.23 | 5/26/99           | 46  |  |

| <b>DOC</b> |          | RMP data |      | Bottom water data |     |  |
|------------|----------|----------|------|-------------------|-----|--|
| Date       | DOC (uM) | DOC (uM) | Date | Sta.              |     |  |
| RC         | 4/22/98  | 177      | 263  | 4/8/98            | 25  |  |
|            |          |          | 267  | 4/8/98            | 46  |  |
| RC         | 7/20/98  | 221      | 341  | 9/16/98           | 25  |  |
| RC&SB      | 4/12/99  | 167      | 141  | 5/26/99           | 29A |  |
|            |          |          | 172  | 5/26/99           | 46  |  |

RC represents RMP's Redwood Creek site

SB represents RMP's San Bruno site

RC&SB is an average of Redwood Creek and San Bruno values

mg/L=milligrams per liter

uM=micromoles per liter

**Table 6** Dissolved Nutrient Flux from Core Incubations on May 26, 1999

| Molar N:P Flux Ratios |      |
|-----------------------|------|
| St. 29A               | 19.0 |
| St. 46                | 25.7 |

| Nitrate (NO3) plus Nitrite              |      |        |                           |                |   |                          | Flux vs. Load comparison |           |            |
|---|------|--------|---------------------------|----------------|---|--------------------------|--------------------------|-----------|------------|
| Nitrate Flux (umoles/m <sup>2</sup> -h) |      |        |                           |                |   |                          | Station*                 | Areally** | City of SJ |
| Station                                 | Core | Mean   | CI                        | r <sup>2</sup> | n | in um/m <sup>2</sup> /hr | in kg/day                | NO3 load  |            |
|   |      |        |                           |                |   | in um/m <sup>2</sup> /hr | in kg/day                | in kg/day |            |
| 29A                                     | 2    | 10 +/- | 18                        | 0.14           | 4 | 32                       | 5873                     | 3831      |            |
| 29A                                     | 3    | 42 +/- | 7                         | 0.95           | 4 | +/-                      | +/-                      |           |            |
| 29A                                     | 4    | 42 +/- | 15                        | 0.79           | 4 | 56                       | 10416                    |           |            |
| 46                                      | 6    | 72 +/- | 97 Initial flux dominated | 0.22           | 4 | 82                       | 15337                    | 3831      |            |
| 46                                      | 7    | 97 +/- | 94 Initial flux dominated | 0.35           | 4 | +/-                      | +/-                      |           |            |
| 46                                      | 8    | 78 +/- | 95 Initial flux dominated | 0.25           | 4 | 40                       | 7439                     |           |            |

| Ammonia (NH4)                           |      |        |                           |                |   |                          | Station*  | Areally** | City of SJ |
|---|------|--------|---------------------------|----------------|---|--------------------------|-----------|-----------|------------|
| Ammonia Flux (umoles/m <sup>2</sup> -h) |      |        |                           |                |   |                          | mean&ci   | averaged  | NH4 load   |
| Station                                 | Core | Mean   | CI                        | r <sup>2</sup> | n | in um/m <sup>2</sup> /hr | in kg/day | in kg/day |            |
| 29A                                     | 2    | 15 +/- | 38                        | 0.07           | 4 | 35                       | 6465      | 487       |            |
| 29A                                     | 3    | 48 +/- | 16                        | 0.81           | 4 | +/-                      | +/-       |           |            |
| 29A                                     | 4    | 41 +/- | 29 Initial flux dominated | 0.51           | 4 | 53                       | 9866      |           |            |
| 46                                      | 6    | 64 +/- | 90 Initial flux dominated | 0.20           | 4 | 72                       | 13334     | 487       |            |
| 46                                      | 7    | 60 +/- | 84 Initial flux dominated | 0.20           | 4 | +/-                      | +/-       |           |            |
| 46                                      | 8    | 92 +/- | 81 Initial flux dominated | 0.39           | 4 | 53                       | 9844      |           |            |

| Reactive Phosphate (DRP)            |      |         |                            |                |   |                          | Station*  | Areally** | City of SJ |
|-------------------------------------|------|---------|----------------------------|----------------|---|--------------------------|-----------|-----------|------------|
| DRP Flux (umoles/m <sup>2</sup> -h) |      |         |                            |                |   |                          | mean&ci   | averaged  | DRP load   |
| Station                             | Core | Mean    | CI                         | r <sup>2</sup> | n | in um/m <sup>2</sup> /hr | in kg/day | in kg/day |            |
| 29A                                 | 2    | 2.3 +/- | 1.5                        | 0.54           | 4 | 3                        | 1439      | 1963      |            |
| 29A                                 | 3    | 2.8 +/- | 0.8                        | 0.86           | 4 | +/-                      | +/-       |           |            |
| 29A                                 | 4    | 5.3 +/- | 1.5                        | 0.86           | 4 | 5                        | 2025      |           |            |
| 46                                  | 6    | 6.8 +/- | 5.3 Initial flux dominated | 0.45           | 4 | 6                        | 2469      | 1963      |            |
| 46                                  | 7    | 6.6 +/- | 5.9 Initial flux dominated | 0.39           | 4 | +/-                      | +/-       |           |            |
| 46                                  | 8    | 4.6 +/- | 6.2 Initial flux dominated | 0.21           | 4 | 4                        | 1540      |           |            |

| Nitrite                                 |      |          |                            |                |   |  |
|---|------|----------|----------------------------|----------------|---|--|
| Nitrite Flux (umoles/m <sup>2</sup> -h) |      |          |                            |                |   |  |
| Station                                 | Core | Mean     | CI                         | r <sup>2</sup> | n |  |
| 29A                                     | 2    | 4.0 +/-  | 3.9 Initial flux dominated | 0.34           | 4 |  |
| 29A                                     | 3    | 12.4 +/- | 1.4                        | 0.97           | 4 |  |
| 29A                                     | 4    | 12.1 +/- | 4.0                        | 0.82           | 4 |  |
| 46                                      | 6    | 5.3 +/-  | 8.1 Initial flux dominated | 0.17           | 4 |  |
| 46                                      | 7    | 6.0 +/-  | 8.2 Initial flux dominated | 0.21           | 4 |  |
| 46                                      | 8    | 6.4 +/-  | 8.1 Initial flux dominated | 0.24           | 4 |  |

| Inorganic Nitroen (DIN)             |      |         |                            |                |   |  |
|-------------------------------------|------|---------|----------------------------|----------------|---|--|
| DIN Flux (umoles/m <sup>2</sup> -h) |      |         |                            |                |   |  |
| Station                             | Core | Mean    | CI                         | r <sup>2</sup> | n |  |
| 29A                                 | 2    | 25 +/-  | 49                         | 0.12           | 4 |  |
| 29A                                 | 3    | 90 +/-  | 20                         | 0.91           | 4 |  |
| 29A                                 | 4    | 83 +/-  | 44                         | 0.64           | 4 |  |
| 46                                  | 6    | 136 +/- | 187 Initial flux dominated | 0.21           | 4 |  |
| 46                                  | 7    | 157 +/- | 178 Initial flux dominated | 0.28           | 4 |  |
| 46                                  | 8    | 169 +/- | 176 Initial flux dominated | 0.32           | 4 |  |

| Silica                                 |      |         |                           |                |   |  |
|--|------|---------|---------------------------|----------------|---|--|
| Silica Flux (umoles/m <sup>2</sup> -h) |      |         |                           |                |   |  |
| Station                                | Core | Mean    | CI                        | r <sup>2</sup> | n |  |
| 29A                                    | 2    | 233 +/- | 69                        | 0.85           | 4 |  |
| 29A                                    | 3    | 242 +/- | 59                        | 0.89           | 4 |  |
| 29A                                    | 4    | 294 +/- | 95                        | 0.83           | 4 |  |
| 46                                     | 6    | 75 +/-  | 40 Initial flux dominated | 0.64           | 4 |  |
| 46                                     | 7    | 90 +/-  | 27                        | 0.85           | 4 |  |
| 46                                     | 8    | 109 +/- | 36                        | 0.82           | 4 |  |

\* Mean of the three flux values for the given station, with 95% confidence interval

\*\* Mean values averaged over the South Bay area (554 km<sup>2</sup>, Cheng and Gartner, 1985)

CI=95% confidence interval

**Table 7** Dissolved Nickel Flux from Core Incubations

| <b>April, 1998</b>                 |      |                  |      |                |    |                                     | Flux vs. Load comparison |                      |            |
|------------------------------------|------|------------------|------|----------------|----|-------------------------------------|--------------------------|----------------------|------------|
| Ni Flux (nmoles/m <sup>2</sup> -h) |      |                  |      |                |    |                                     | Station*                 | Areally**            | City of SJ |
| Station                            | Core | Mean             | CI   | r <sup>2</sup> | n  | mean&ci<br>in nM/m <sup>2</sup> /hr | averaged<br>in kg/day    | Ni load<br>in kg/day |            |
| 25                                 | 6    | <b>25.9</b> +/-  | 4.3  | 0.48           | 17 | 14.0                                | <b>10.9</b>              | <b>3.0</b>           |            |
| 25                                 | 7    | <b>14.5</b> +/-  | 8.4  | 0.16           | 12 | +/-                                 | +/-                      |                      |            |
| 25                                 | 8    | <b>1.7</b> +/-   | 2.5  | 0.02           | 13 | 36.8                                | 28.8                     |                      |            |
| 46                                 | 1    | <b>65.1</b> +/-  | 7.7  | 0.72           | 15 | 73.3                                | <b>57.2</b>              | <b>3.0</b>           |            |
| 46                                 | 3    | <b>116.3</b> +/- | 10.5 | 0.81           | 15 | +/-                                 | +/-                      |                      |            |
| 46                                 | 4    | <b>38.6</b> +/-  | 2.8  | 0.95           | 10 | 120.3                               | 93.9                     |                      |            |

| <b>September, 1998</b>             |      |                  |      |                |    |                                     | Flux vs. Load comparison |                      |            |
|------------------------------------|------|------------------|------|----------------|----|-------------------------------------|--------------------------|----------------------|------------|
| Ni Flux (nmoles/m <sup>2</sup> -h) |      |                  |      |                |    |                                     | Station*                 | Areally**            | City of SJ |
| Station                            | Core | Mean             | CI   | r <sup>2</sup> | n  | mean&ci<br>in nM/m <sup>2</sup> /hr | averaged<br>in kg/day    | Ni load<br>in kg/day |            |
| 25                                 | 1    | <b>94.6</b> +/-  | 8.3  | 0.79           | 16 | 55.4                                | <b>43.2</b>              | <b>3.0</b>           |            |
| 25                                 | 2    | <b>32.3</b> +/-  | 8.2  | 0.44           | 13 | +/-                                 | +/-                      |                      |            |
| 25                                 | 3    | <b>106.3</b> +/- | 19.2 | 0.61           | 13 | 101.5                               | 79.2                     |                      |            |
| 25                                 | 4    | <b>-11.6</b> +/- | 13.9 | 0.02           | 15 |                                     |                          |                      |            |

| <b>May, 1999</b>                   |      |                 |      |                |    |                                     | Flux vs. Load comparison |                      |            |
|------------------------------------|------|-----------------|------|----------------|----|-------------------------------------|--------------------------|----------------------|------------|
| Ni Flux (nmoles/m <sup>2</sup> -h) |      |                 |      |                |    |                                     | Station*                 | Areally**            | City of SJ |
| Station                            | Core | Mean            | CI   | r <sup>2</sup> | n  | mean&ci<br>in nM/m <sup>2</sup> /hr | averaged<br>in kg/day    | Ni load<br>in kg/day |            |
| 29A                                | 2    | <b>39.9</b> +/- | 7.4  | 0.60           | 13 | 64.5                                | <b>50.3</b>              | <b>3.0</b>           |            |
| 29A                                | 3    | <b>66.1</b> +/- | 9.2  | 0.72           | 13 | +/-                                 | +/-                      |                      |            |
| 29A                                | 4    | <b>87.4</b> +/- | 12.4 | 0.64           | 15 | 72.5                                | 56.6                     |                      |            |
| 46                                 | 6    | <b>53.8</b> +/- | 5.3  | 0.79           | 15 | 60.5                                | <b>47.2</b>              | <b>3.0</b>           |            |
| 46                                 | 7    | <b>33.7</b> +/- | 4.6  | 0.70           | 14 | +/-                                 | +/-                      |                      |            |
| 46                                 | 8    | <b>94.1</b> +/- | 12.1 | 0.76           | 13 | 93.7                                | 73.1                     |                      |            |

\* Mean of the three (or four) flux values for the given station, with 95% confidence interval

\*\* Mean values averaged over the South Bay area (554 km<sup>2</sup>, Cheng and Gartner, 1985)

CI=95% confidence interval

| Average and CI for Ni flux estimates in kg/day |     |             |      |
|--|-----|-------------|------|
| <b>41.9</b>                                    | +/- | <b>16.4</b> | n=16 |

**Table 8.** Comparison of nutrient fluxes using two or twelve hours of incubation data

| All dissolved fluxes        |   |             |
|-----------------------------|---|-------------|
| St.46                       | in $\mu\text{mol}/\text{m}^2/\text{hr}$ |             |
| Core                        | All 12 hrs                              | First 2 hrs |
| <b>Nitrite</b>              |   |             |
| 6                           | 5                                       | 74          |
| 7                           | 6                                       | 76          |
| 8                           | 6                                       | 78          |
| <b>Nitrate plus nitrite</b> |   |             |
| 6                           | 72                                      | 890         |
| 7                           | 97                                      | 904         |
| 8                           | 78                                      | 882         |
| <b>Phosphate</b>            |   |             |
| 6                           | 7                                       | 53          |
| 7                           | 7                                       | 57          |
| 8                           | 5                                       | 58          |
| <b>Silica</b>               |   |             |
| 6                           | 75                                      | 395         |
| 7                           | 90                                      | 240         |
| 8                           | 109                                     | 433         |
| <b>Ammonia</b>              |   |             |
| 6                           | 64                                      | 836         |
| 7                           | 60                                      | 789         |
| 8                           | 92                                      | 807         |
| <b>Inorganic nitrogen</b>   |   |             |
| 6                           | 136                                     | 1726        |
| 7                           | 157                                     | 1693        |
| 8                           | 169                                     | 1690        |

**Table 9.** Dissolved Organic Carbon Flux from Core Incubations

| <b>April, 1998</b> |      | DOC Flux (umoles/m <sup>2</sup> -h) |     |                |    |  |
|--------------------|------|-------------------------------------|-----|----------------|----|--|
| Station            | Core | Mean                                | CI  | r <sup>2</sup> | n  |  |
| 25                 | 6    | <b>539</b> +/-                      | 92  | 0.59           | 14 |  |
| 25                 | 7    | <b>-32</b> +/-                      | 38  | 0.02           | 18 |  |
| 25                 | 8    | <b>-71</b> +/-                      | 14  | 0.42           | 16 |  |
| 46                 | 1    | <b>676</b> +/-                      | 160 | 0.47           | 13 |  |
| 46                 | 3    | <b>-222</b> +/-                     | 80  | 0.24           | 14 |  |
| 46                 | 4    | <b>89</b> +/-                       | 17  | 0.48           | 15 |  |

| <b>September, 1998</b> |      | DOC Flux (umoles/m <sup>2</sup> -h) |    |                |    |  |
|------------------------|------|-------------------------------------|----|----------------|----|--|
| Station                | Core | Mean                                | CI | r <sup>2</sup> | n  |  |
| 25                     | 1    | <b>328</b> +/-                      | 58 | 0.62           | 13 |  |
| 25                     | 2    | <b>-68</b> +/-                      | 55 | 0.06           | 14 |  |
| 25                     | 3    | <b>-80</b> +/-                      | 12 | 0.70           | 13 |  |
| 25                     | 4    | <b>-18</b> +/-                      | 27 | 0.02           | 13 |  |

| <b>May, 1999</b> |      | DOC Flux (umoles/m <sup>2</sup> -h) |    |                |    |  |
|------------------|------|-------------------------------------|----|----------------|----|--|
| Station          | Core | Mean                                | CI | r <sup>2</sup> | n  |  |
| 29A              | 2    | <b>59</b> +/-                       | 24 | 0.17           | 15 |  |
| 29A              | 3    | <b>103</b> +/-                      | 19 | 0.54           | 14 |  |
| 29A              | 4    | <b>24</b> +/-                       | 28 | 0.02           | 15 |  |
| 46               | 6    | <b>-239</b> +/-                     | 47 | 0.40           | 17 |  |
| 46               | 7    | <b>-96</b> +/-                      | 53 | 0.12           | 14 |  |
| 46               | 8    | <b>-85</b> +/-                      | 55 | 0.11           | 13 |  |

CI=95% confidence interval

**Table 10.** Dissolved Sulfide Flux from Core Incubations

| <b>April, 1998</b> |      | Sulfide Flux (nmoles/m <sup>2</sup> -h) |     |     |                |   |
|--------------------|------|---|-----|-----|----------------|---|
| Station            | Core | Mean                                    |     | CI  | r <sup>2</sup> | n |
| 25                 | 6    | <b>752</b>                              | +/- | 103 | 0.95           | 7 |
| 25                 | 7    | <b>648</b>                              | +/- | 52  | 0.97           | 8 |
| 25                 | 8    | <b>647</b>                              | +/- | 71  | 0.95           | 8 |
| 46                 | 1    | <b>499</b>                              | +/- | 41  | 0.97           | 8 |
| 46                 | 3    | <b>638</b>                              | +/- | 57  | 0.96           | 8 |
| 46                 | 4    | <b>574</b>                              | +/- | 52  | 0.96           | 8 |

| <b>September, 1998</b> |      | Sulfide Flux (nmoles/m <sup>2</sup> -h) |     |    |                |   |
|------------------------|------|---|-----|----|----------------|---|
| Station                | Core | Mean                                    |     | CI | r <sup>2</sup> | n |
| 25                     | 1    | <b>345</b>                              | +/- | 40 | 0.94           | 8 |
| 25                     | 2    | <b>458</b>                              | +/- | 39 | 0.97           | 8 |
| 25                     | 3    | <b>480</b>                              | +/- | 21 | 0.99           | 8 |
| 25                     | 4    | <b>398</b>                              | +/- | 37 | 0.96           | 8 |

| <b>May, 1999</b> |      | Sulfide Flux (nmoles/m <sup>2</sup> -h) |     |    |                |   |
|------------------|------|---|-----|----|----------------|---|
| Station          | Core | Mean                                    |     | CI | r <sup>2</sup> | n |
| 29A              | 2    | <b>325</b>                              | +/- | 25 | 0.97           | 8 |
| 29A              | 3    | <b>312</b>                              | +/- | 8  | 0.99           | 8 |
| 29A              | 4    | <b>357</b>                              | +/- | 27 | 0.98           | 7 |
| 46               | 6    | <b>183</b>                              | +/- | 40 | 0.81           | 8 |
| 46               | 7    | <b>211</b>                              | +/- | 32 | 0.94           | 7 |
| 46               | 8    | <b>181</b>                              | +/- | 15 | 0.98           | 7 |

CI=95% confidence interval