

NOAA Technical Memorandum NOS ORCA 100



Contaminants in Aquatic Habitats at Hazardous Waste Sites: Mercury

December 1996
Seattle, Washington

noaa NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION

National Ocean Service

**Office of Ocean Resources Conservation and Assessment
National Ocean Service
National Oceanic and Atmospheric Administration
U.S. Department of Commerce**

The Office of Ocean Resources Conservation and Assessment (ORCA) provides decisionmakers comprehensive, scientific information on characteristics of the oceans, coastal areas, and estuaries of the United States of America. The information ranges from strategic, national assessments of coastal and estuarine environmental quality to real-time information for navigation or hazardous materials spill response. Through its National Status and Trends (NS&T) Program, ORCA uses uniform techniques to monitor toxic chemical contamination of bottom-feeding fish, mussels and oysters, and sediments at about 300 locations throughout the United States. A related NS&T Program of directed research examines the relationships between contaminant exposure and indicators of biological responses in fish and shellfish.

Through the Hazardous Materials Response and Assessment Division (HAZMAT) Scientific Support Coordination program, ORCA provides critical scientific support to the U.S. Coast Guard for planning and responding to spills of oil or hazardous materials into marine or estuarine environments. Technical guidance includes spill trajectory predictions, chemical hazard analyses, and assessments of the sensitivity of marine and estuarine environments to spills. To fulfill the responsibilities of the Secretary of Commerce as a trustee for living marine resources, HAZMAT's Coastal Resource Coordination program provides technical support to the U.S. Environmental Protection Agency during all phases of the remedial process to protect the environment and restore natural resources at hundreds of waste sites each year. As another part of its marine trusteeship responsibilities, ORCA conducts comprehensive assessments of damages to coastal and marine resources from discharges of oil and hazardous materials.

ORCA collects, synthesizes, and distributes information on the use of the coastal and oceanic resources of the United States to identify compatibilities and conflicts and to determine research needs and priorities. It conducts comprehensive, strategic assessments of multiple resource uses in coastal, estuarine, and oceanic areas for decisionmaking by NOAA, other Federal agencies, state agencies, Congress, industry, and public interest groups. It publishes a series of thematic data atlases on major regions of the U.S. Exclusive Economic Zone and on selected characteristics of major U.S. estuaries.

ORCA implements NOAA responsibilities under Title II of the Marine Protection, Research, and Sanctuaries Act of 1972; Section 6 of the National Ocean Pollution Planning Act of 1978; the Oil Pollution Act of 1990; the National Coastal Monitoring Act of 1992; and other Federal laws. It has four major line organizations: Coastal Monitoring and Bioeffects Assessment Division, Hazardous Materials Response and Assessment Division, Strategic Environmental Assessment Division, and the Damage Assessment Center.

NOAA Technical Memorandum NOS ORCA 100

Contaminants in Aquatic Habitats at Hazardous Waste Sites: Mercury

Nancy Beckvar ¹
Jay Field ¹
Sandra Salazar ²
Rebecca Hoff ¹

¹NOAA/HAZMAT
²EVS Consultants
Seattle, Washington



Seattle, Washington

United States
Department of Commerce
Mickey Kantor
Secretary

National Oceanic and
Atmospheric Administration
D. James Baker
Under Secretary for Oceans
and Atmosphere

National Ocean Service
W. Stanley Wilson
Assistant Administrator
for Ocean Services and
Coastal Zone Management

**Hazardous Materials Response and Assessment Division
Office of Ocean Resources Conservation and Assessment
National Ocean Service
National Oceanic and Atmospheric Administration
U.S. Department of Commerce
Silver Spring, Maryland**

CITATION

Please cite this report as "Beckvar, Nancy, Jay Field, Sandra Salazar, and Rebecca Hoff. 1996. *Contaminants in Aquatic Habitats at Hazardous Waste Sites: Mercury*. NOAA Technical Memorandum NOS ORCA 100. Seattle: Hazardous Materials Response and Assessment Division, National Oceanic and Atmospheric Administration. 74 pp.

NOTICE

This report has been reviewed by the National Ocean Service of the National Oceanic and Atmospheric Administration (NOAA) and approved for publication. Such approval does not signify that the contents of this report necessarily represent the official position of NOAA or of the Government of the United States, nor does mention of trade names or commercial products constitute endorsement or recommendation for their use.

Contents

Executive Summary	i
Environmental Chemistry	i
Bioaccumulation	ii
Toxicity	iii
Applications	iv
Summary	vi
Acronyms	vii
Glossary	viii
Introduction	1
Environmental Chemistry	1
Chemical Speciation	2
Distribution in the Environment	9
Bioaccumulation of Mercury	12
The Effect of the Form of Mercury on Bioaccumulation	13
Exposure Pathways	17
Biological Factors Affecting Accumulation of Mercury	21
Other Factors Affecting Accumulation	23
Toxicity of Mercury	25
Toxicity of Mercury in Water	27
Toxicity of Mercury in Sediment	32
Toxicity Associated with Mercury in Tissues	33
Interactions with Other Metals	36
Criteria and Guidelines	36
Ambient Water Quality Criteria	36
Sediment Guidelines	38
Tissue	38

Contents, cont.

Applications	38
Sampling and Monitoring Considerations	39
Approaches to Remediation	41
Analytical Considerations.....	43
Summary	44
References.....	45
Figure	
1 Mercury speciation.....	3
Tables	
1 Toxicity of Mercury in Water.....	28
2 Toxicity Associated with Mercury in Tissues.....	35

EXECUTIVE SUMMARY

This document reviews published literature on mercury chemistry, bioaccumulation and toxicity, and is intended to serve as guidance for NOAA Coastal Resource Coordinators in their work with EPA on hazardous waste sites. The purpose of this document is to highlight factors to consider in designing and evaluating ecological risk assessments; and, in the sampling, monitoring and analyses of environmental media in aquatic habitats affected by mercury. Though many questions about mercury remain, the science is evolving rapidly. This paper should be reviewed with the knowledge that information can change as new studies are published.

Environmental Chemistry

The fate of mercury in the environment depends on the chemical form of mercury released and the environmental conditions. Elemental mercury, inorganic mercury, and methylmercury are the three most important forms of mercury in natural aquatic environments. Most mercury is released into the environment as inorganic mercury, which is primarily bound to particulates and organic substances and may not be available for direct uptake by aquatic organisms. The process of methylation of inorganic mercury to methylmercury, which is highly bioavailable, is thus an important key to the fate of mercury in the environment.

Elemental mercury has a high vapor pressure, a low solubility, does not combine with inorganic or organic ligands, and is not available for methylation. The mercurous ion (Hg[I]) combines with inorganic compounds only and cannot be methylated. The mercuric ion (Hg[II]) combines with both inorganic and organic ligands, and can be methylated. Methylation in aquatic habitats is primarily a biological process. Mono- and dimethylmercury are formed by microorganisms in both sediment and water through the methylation of inorganic mercuric ions (Hg[II]). Dimethylmercury, which is highly volatile, is generally not persistent in aquatic environments.

Methylation is influenced by environmental variables that affect both the availability of mercuric ions for methylation and the growth of the methylating microbial populations. Methylation rates are higher under anoxic conditions, in

freshwater compared to saltwater, and in low pH environments. The presence of organic matter can stimulate growth of microbial populations (and reduce oxygen levels), thereby enhancing the formation of methylmercury. Sulfide can bind mercury and limit methylation. Methylmercury production can vary due to seasonal changes in nutrients, oxygen, temperature, and hydrodynamics. In most studies, methylation increased during the summer months when biological productivity was high, and decreased during the winter months.

Measurements of total mercury concentrations in the sediment do not provide information on the form of mercury present, methylation potential, or availability to organisms locally and downstream. If environmental conditions are conducive for methylation, methylmercury concentrations may be high in proportion to the supply and distribution of total mercury.

Bioaccumulation

Mercury is accumulated by fish, invertebrates, mammals, and aquatic plants and the concentration tends to increase with increasing trophic level (mercury biomagnifies). Although inorganic mercury is the dominant form of mercury in the environment and is easily taken up, it is also depurated relatively quickly. Methylmercury accumulates quickly, depurates very slowly, and therefore biomagnifies in higher trophic species. The percentage of methylmercury, as compared to total mercury, also increases with age in both fish and invertebrates. Uptake and depuration rates vary between tissues within an organism. Partitioning of mercury between tissues within aquatic organisms is influenced by the chemical form of mercury and route of exposure (ingestion or via the gills). Due to its preferential uptake, ability to be transferred among tissues, and slow depuration, most of the mercury in fish muscle tissue (99%) is methylmercury.

Marine mammal tissues have some of the highest concentrations of mercury found in all marine organisms, with the liver generally having the highest total mercury concentration. Although many juvenile and adult marine mammals primarily feed on fish, which contain high percentages of methylmercury, high concentrations of inorganic mercury are found in adult specimens. Apparently, adult marine mammals can mineralize methylmercury into inorganic mercury. Juvenile marine mammals have lower concentrations of total mercury than adults; but unlike fish and invertebrates, the percentage of methylmercury is higher in juvenile mammals.

Invertebrates generally have a lower percentage of methylmercury, as compared to total mercury, in their tissues than do fish and marine mammals. The percentage of methylmercury in invertebrates varies greatly and can range from one percent in deposit-feeding polychaetes, to close to 100% in crab.

Bioconcentration factors (BCFs) reflect uptake from water in laboratory experiments. BCFs for mercury are variable, with the highest factors determined for methylmercury. BCFs for methylmercury in brook trout range from 69,000 to 630,000, depending on the tissue analyzed. BCFs for inorganic mercury (mercuric chloride) in saltwater species range from 129 for adult lobster (*Homarus americanus*) to 10,000 for oysters (*Crassostrea virginica*).

While sediment is usually the primary source of mercury in most aquatic systems, the food web is the main pathway for accumulation. High trophic level species tend to accumulate the highest concentrations of mercury, with concentrations highest in fish-eating predators. Mercury concentrations in higher trophic species often do not correlate with concentrations in environmental media. Correlations have been made between sediment and lower trophic species that typically have a high percentage of inorganic mercury, and between mercury concentrations in higher trophic species and their prey items. The best measure of bioavailability of mercury in any system can be obtained by analyzing mercury concentrations in the biota at the specific site.

Toxicity

Toxicity is influenced by the form of mercury, the environmental media, environmental conditions, the sensitivity or tolerance of the organism, and the life history stage. Inorganic mercury is less acutely toxic to aquatic organisms than methylmercury, but the range in sensitivity among individual species for either compound is large. Toxicity was found to be greater at elevated temperatures, lower oxygen content, reduced salinities in marine environments, and in the presence of metals such as zinc and lead.

In general, toxic effects occur because mercury binds to proteins and alters protein production or synthesis. Toxicological effects include reproductive impairment, growth inhibition, developmental abnormalities, and altered behavioral responses.

Reproductive endpoints are generally more sensitive than growth or survival, with embryos and the early developmental stages the most sensitive. Mercury can be transferred from tissues of the adult female to developing eggs. Exposure to low concentrations of mercury may not result in mortality directly, but may retard growth thereby increasing the risk of predation.

Data available on the effects of mercury-contaminated sediment on aquatic organisms reviewed by Long and MacDonald (1992) resulted in effects range-low (ERL) and effects range-median (ERM) concentrations of 0.15 mg/kg and 0.71 mg/kg, respectively. However, these numbers were less accurate than other metals in predicting adverse effects, highlighting the need for site-specific effects data to determine concentrations of mercury in sediment that pose a threat to aquatic biota.

Few studies report both tissue residues and effects in long-term exposure to low concentrations of mercury. However, results from studies on different freshwater species indicate that reproductive effects could be expected to occur in sensitive fish species at tissue concentrations close to the FDA action level of 1 mg/kg (ppm). The interaction of mercury and other trace elements (e.g., cadmium, copper, selenium, and zinc) can be both antagonistic and synergistic, primarily depending on exposure concentrations and form of mercury. Effects were generally less than additive (antagonistic) at lower exposure levels and greater than additive (synergistic) at higher levels. Zinc and cadmium were reported to reduce the teratogenic effects of methylmercury to killifish while selenium reduced mercury's toxic effects on development in medaka embryos.

Applications

Ecological assessments of waste sites with elevated concentrations of mercury in the aquatic environment are particularly challenging due to the complexity of the factors that affect the availability of mercury to aquatic organisms. Depending on the magnitude of the problem (local versus system-wide), the level of effort necessary to evaluate mercury contamination may range from simple monitoring of chemical concentrations to more complex programs including monitoring of numerous physical, chemical and biological parameters. The distribution of total mercury in sediment, which in most cases is predominantly inorganic mercury, may not by itself provide useful information about the bioavailability of mercury to

aquatic species. Concentrations of total mercury in sediment that decrease with increasing distance from the source may still pose a threat to organisms if the bioavailability of the mercury increases (i.e., environmental conditions are more conducive for methylation). Mercury concentrations in aquatic organisms, particularly higher trophic-level organisms, may provide the best measure of the availability of mercury in a particular area.

In sites where a whole system has been affected, evaluation of remedial alternatives may need to be based on an understanding of the system-specific processes that lead to increased methylation and the pathways to resources of concern. An assessment of environmental parameters that affect the activity of methylating microbes (e.g., nutrients, temperature, pH, and dissolved oxygen) and the factors affecting the availability of inorganic mercury for methylation (e.g., the resuspension of sediment, total organic carbon, and sulfides) may be warranted when designing sampling plans for a remedial investigation.

To establish protective sediment target cleanup concentrations and remedial options for mercury-contaminated sites, we must understand the extent of contamination, the major pathways of transport, and bioavailability. Therefore, data on the accumulation of mercury in tissues of aquatic organisms should be included in assessment studies because it addresses potential human health concerns and availability to aquatic receptors. In addition, studies should assess toxicity to aquatic organisms, focusing on early life stages and reproductive effects.

Detection limits should reflect the program objectives. Contract Lab Program (CLP) methodology may be appropriate for screening level assessments; however, biologically relevant detection limits are often required and not available at CLP laboratories. Thus, analytical laboratories that can achieve lower detection limits may need to be used. Quality control is an important aspect of any testing program but is particularly important when analyzing mercury in environmental matrices. In water, very low concentrations need to be measured; the separation of the different forms of mercury requires special analytical techniques. Matrix effects in the extraction of mercury from tissue may interfere with accurate analyses for methyl and total mercury. When analyzing mercury in water, sediment, and tissue, analysis of certified standards for the appropriate matrix must be included as part of the quality control plan.

Summary

NOAA recommends a site-specific approach that focuses on determining the availability of mercury and the potential for toxic effects. The accumulation of mercury in aquatic biota is often the primary concern at mercury sites and is useful for assessing availability. Bioaccumulation studies should measure tissue concentrations in more than one resident and/or transplanted caged species, preferably with species representing different trophic levels or different food web pathways. It may not be possible to correlate sediment mercury concentrations with concentrations in biota. However, correlations between mercury concentrations in predator and prey species may be useful in determining pathways of mercury transfer.

Toxicity tests such as the standard amphipod tests should also be conducted to assess mercury toxicity to benthic organisms. At major mercury sites, chronic toxicity endpoints should be included in the assessment—in particular, fish early life stage or reproductive endpoint tests. Because of the persistence of mercury in aquatic systems, source control alone may not be sufficient to permit recovery. Additional remedial actions may be required to reduce the total mercury burden in the system. Long-term monitoring of tissue concentrations of mercury in aquatic biota is needed to assess remedial effectiveness.

ACRONYMS AND ABBREVIATIONS

AVS	acid-volatile sulfides
AWQC	Ambient Water Quality Criteria
BAFs	Bioaccumulation factors
BCFs	Bioconcentration factors
Cd	cadmium
CLP	Contract Lab Program
Cu	copper
DOC	dissolved organic carbon
Eh	oxidation-reduction potential
ERL	effects range-low
ERM	effects range-median
FDA	U.S. Food and Drug Administration
Fe	iron
Hg	mercury
Mn	manganese
MT	metallothioneins
$\mu\text{eq/l}$	micro equivalent per liter
$\mu\text{g/l}$	micrograms per liter
μm	micrometer
mg/kg	milligrams per kilogram
mg/l	milligrams per liter
ppm	parts per million
Se	selenium
SRB	sulfate-reducing bacteria
TOC	total organic carbon
Zn	zinc

GLOSSARY

acid labile mercury

Determined by SnCl_2 reduction on acidified samples, includes inorganic complexes, labile organic associations, elemental mercury, and labile particulate mercury. Doesn't measure organic forms (C-Hg bound) of mercury such as methylmercury. Same as reactive mercury.

acid soluble mercury

Mercury that passes through a 0.45 μm membrane filter after the sample is acidified to pH 1-2.0 with nitric acid (EPA 1984). Strongly sorbed Hg is not measured, but all toxic forms as well as some non-toxic forms are measured.

alkylmercury

Includes phenyl-, monomethyl-, and dimethylmercury

bioaccumulation

Net uptake of a contaminant into tissue from all pathways

bioaccumulation factors (BAF)

Ratio of tissue concentration to concentration in medium, with exposure from the food web and the medium

bioconcentration factors (BCF)

Ratio of tissue concentration to concentration in medium, with exposure only through the medium

biomagnification

Tissue concentration increases as trophic level increases

DOC (dissolved organic carbon)

Includes all sources of carbon, including humic and fulvic matter as well as carbohydrates, proteins, etc.

demethylation

Conversion of methylmercury back to an inorganic form.

depuration

Elimination of a contaminant from the body of an organism

dimethylmercury

Organic form of mercury consisting of a single mercury atom and two methyl groups $[(\text{CH}_3)_2\text{Hg}]$, highly volatile and not persistent in the environment.

Effects range-low

Concentration equivalent to that reported at the lower 10 percentile of the

available screened sediment toxicity data for predominantly marine and estuarine toxicity studies. This number represents the low end of the range of concentrations at which effects were observed in the studies compiled (Long and MacDonald 1992).

Effects range-median

Concentration equivalent to that reported at the 50th percentile of the available screened sediment toxicity data for predominantly marine and estuarine toxicity studies (Long and MacDonald 1992).

elemental mercury

Not in ionic form, cannot be methylated

halide

Binary compound of a halogen (fluorine, chlorine, bromide, iodine and astatine)

inorganic mercury

Includes elemental mercury and mercury bound to other inorganic molecules and compounds, including inorganic ligands and sulfides.

labile mercury

Includes $\text{Hg}(\text{OH})_2$, HOHgCl , HgCl_2 , and weakly bound organo-complex forms

ligand

Any group, ion, or molecule that binds to another, called a receptor

methylmercury

Includes both mono- and dimethylmercury

mercuric-Hg[II]

Forms both inorganic and organic complexes, this is the only ionic form that can be methylated

mercurous-Hg[I]

Forms inorganic complexes, cannot be methylated

methylation

Addition of a methyl (CH_3) group

monomethylmercury

Organic form of mercury with one methyl group attached to a mercury atom (CH_3Hg) - highly toxic and readily accumulated by living organisms

organic mercury

Includes mercury complexes with organic ligands (e.g. humic/fulvic acids,

amino acids, but without a Hg-carbon bond) and organic mercury bound via a carbon atom ($\text{CH}_3\text{-HgOH}$, CH_3HgCl , CH_3HgCH_3)

organo-mercury

Mercury compounds with a Hg-carbon bond

pinniped

Mammals of the family Pinnipedia, includes all seals and walruses

piscivorous

Feeds on fishes

reactive mercury

“Easily reducible,” determined by SnCl_2 reduction on acidified samples, includes inorganic complexes, labile organic associations, elemental mercury, and labile particulate mercury, doesn't include C-Hg bound mercury such as methylmercury and dimethylmercury. Same as acid labile mercury.

total mercury

Includes all forms of mercury

INTRODUCTION

As a trustee for marine, estuarine, and anadromous resources, NOAA is responsible for ensuring the well-being of those trust resources potentially affected by releases from hazardous waste sites. Metals, in particular, pose a threat because of their persistence and toxicity in aquatic environments. The importance of mercury in many aquatic environments is underscored by the fact that 35 states have fish and wildlife consumption advisories in place for mercury (EPA 1996).

Although the hazards of mercury to humans are well-known, less information is available on the risks to aquatic organisms. In order to define and address the potential effects and extent of mercury contamination at hazardous waste sites, a number of questions are often asked: Is the mercury present in a bioavailable form? What concentrations in water, sediment, and tissues are potentially harmful to aquatic resources? What types of sampling and analysis are necessary to define potential risks to aquatic organisms? Is there a relationship between sediment concentrations and tissue concentrations in aquatic organisms? What level should be used for cleanup? What are important factors to consider when selecting a remedy and designing monitoring plans at a site?

ENVIRONMENTAL CHEMISTRY

Mercury is among the most toxic of the heavy metals, has complex behavior in the environment, and may persist for decades following abatement of the source. Mercury's environmental persistence is due in part to its high affinity for particulates and organic matter. Even if mercury concentrations in sediment and water decrease over time, concentrations in organisms may not decrease due to the slow rate of elimination of the highly bioavailable methylmercury form. The physical properties, bioavailability, and toxicity of mercury are governed by speciation into both organic and inorganic forms.

Elemental mercury, bivalent inorganic mercury, and monomethylmercury are the three most important forms of mercury occurring in natural aquatic environments (Battelle 1987). Elemental mercury in aquatic environments has a high vapor pressure, a low solubility in water, and an octanol-water partition coefficient (K_{OW})=4.15 (Shoichi and Sokichi 1985 as cited in Major et al. 1991). Elemental and

dimethylmercury can occur as dissolved gaseous mercury. Mercury can also occur as particulate and dissolved ionic and monomethylmercury species. In natural water, ionic mercury is consumed by methylation, reduction, and particulate scavenging (Mason et al. 1995a). Bivalent inorganic mercury binds to inorganic and organic ligands, especially sulfur-containing ligands, and forms both inorganic and organic complexes. Figure 1 shows a schematic of some common pathways of mercury speciation in the environment.

Although most mercury occurs in the inorganic form, methylmercury, an organic form, is the most toxic and readily bioaccumulated form of mercury. Methylmercury normally occurs in the environment at extremely low concentrations; however, it is taken up easily by aquatic organisms and bioaccumulated. Consequently, methylmercury may comprise more than 95% of the mercury in fish tissue while only 5-15% of the total mercury burden in sediments and water of contaminated lakes is methylmercury (Saroff 1990).

Chemical Speciation

Inorganic and total mercury versus methylmercury

Chemical speciation terms commonly used include total, inorganic, organic, and methylmercury and are based on the oxidation state and associated compounds. Mercury has three stable oxidation states: the native element (Hg[0]), mercurous (Hg[I]), and mercuric (Hg[II]). Inorganic mercury includes elemental Hg and some complexes of the mercurous and mercuric oxidation states. Hg [I] forms inorganic compounds only and, like Hg[0], cannot be methylated. Hg [I] compounds include mercurous salts (halides) and mercurous chlorides such as calomel. Both Hg[0] and Hg[I] can be oxidized to form Hg[II]. Hg[II] (bivalent mercury), the form that can be methylated, forms both organic and inorganic compounds. Mercury [II] can combine with inorganic ligands including chloride, hydroxide, nitrate, and sulfate anions (Benes and Havlik 1979) to form inorganic mercury compounds that include mercuric halides, mercuric chloride (cinnabar), and mercuric sulfides. Chloride concentration and pH affect the proportions of the uncharged inorganic species in solution. For example, at low chloride concentrations most of the inorganic mercury occurs in the form of mercuric hydroxide (Hg(OH)₂), with mercuric chloride (HgCl₂) and mercury hydroxide chloride (HgOHCl) also important (Mason et al. 1996). As chloride concentration increases




Figure 1. Important pathways of mercury speciation in the aquatic environment.

(e.g., high-chloride lake water), the proportion of HgCl_2 increases and the other two species constitute only a few percent of the total inorganic mercury. As pH increases, more Cl is needed than at lower pHs to increase the percent of HgCl_2 . At even higher chloride concentrations, HgCl_4^{2-} becomes the dominant species. This speciation chemistry affects the accumulation and toxicity as described later in this report.

The term organic mercury can include different types of organically bound mercury. Hg [II] combines with organic compounds (humic/fulvic acids, amino acids) via an organic ligand bond to form organomercury salts. The Hg [II]-organic ligand bond is relatively weak compared to a C-Hg bond. The organomercury salts resemble their corresponding inorganic mercuric salts in their properties and reactions. The organomercury compounds methylmercury, dimethylmercury, and phenylmercury have a C-Hg bond; methyl (CH_3) and phenyl groups (C_6H_5) link to a mercury atom via a carbon atom. Some authors group the organically bound Hg [II] complexes (without a C-Hg bond) with inorganic mercury compounds, while others group all organically bound mercury together as organic mercury.

Mercury compounds may be grouped according to their form based on chemical speciation discussed above, or based on the analytical technique used to measure the mercury. The analytical technique determines which forms of mercury are detected. Mercury terms based on analytical procedures include names such as acid-soluble, reactive or acid labile, and calcium chloride-extractable.

Measurements of reactive mercury include Hg [II] bound to inorganic substances and weakly bound to organic matter (however, methylmercury is not included due to the strong C-Hg bond). To estimate methylmercury concentrations, some authors measure total mercury and reactive mercury, then subtract the reactive from the total measurement. Total mercury measurements include all the various inorganic and organic forms of mercury, including dissolved, colloidal, and/or particulate states. Analytical groupings are defined in the glossary. Unless the specific type of mercury is mentioned, the reader should assume that the term "mercury" refers to total mercury.

Speciation of mercury is affected by environmental conditions such as pH, oxidation-reduction potential (Eh), oxygen content, sulfide content, chloride

concentration, organic matter content, and microbial activity. Similarly, biological and chemical processes control the conversion of inorganic mercury to methylmercury. The factors that enhance and inhibit methylation, and affect the distribution of inorganic and methylmercury are discussed in this section of the paper.

Methylation of Mercury

In both freshwater and saltwater environments, mercury is converted from inorganic bivalent mercury (Hg[II]) to methylmercury primarily by microorganisms (Berman and Bartha 1986), although chemical methylation also occurs (Craig and Moreton 1985; Weber 1993). Two forms of monomethylmercury, methylmercuric hydroxide (CH_3HgOH) and methylmercuric chloride (CH_3HgCl) occur in both fresh and saltwater, with the former dominant at low chloride concentrations (low chloride freshwater) and the latter dominant at high chloride concentrations (high chloride lakes, seawater). As with inorganic mercury, the organic chloride species ($K_{\text{OW}} = 1.7$) is more hydrophobic than the hydroxide species ($K_{\text{OW}} = 0.07$; Major et al. 1991; Faust 1992; Mason et al. 1996). Dimethylmercury ($K_{\text{OW}} = 182$) readily volatilizes from surface water and is generally not persistent in aquatic environments at concentrations of concern; therefore, discussions of methylmercury in this review refer to the monomethylmercury species, unless otherwise stated.

Methylmercury production depends on both the availability of Hg[II] for methylation and microbial activity. Methylation is usually greatest at the sediment-water interface, but also occurs in the water column. Net methylmercury production is a function of both the rate of methylation and the rate of demethylation (Korthals and Winfrey 1987). Methylmercury is not readily decomposed so the methylation rate is usually higher than the demethylation rate. Degradation of methylmercury is also primarily a microbial process.

Methylation is influenced by the availability of Hg[II], oxygen concentration, pH, redox potential (Eh), presence of sulfate and sulfide, type and concentrations of complexing inorganic and organic agents (Parks et al. 1989), salinity (Blum and Bartha 1980), and organic carbon (Jackson 1989; Winfrey and Rudd 1990). Strongly bound Hg[II] is not available for methylation. For example, insoluble mercuric

sulfide (HgS) will be methylated in aerobic sediments at rates 100 to 1,000 times slower than for the less strongly bound HgCl₂ (Olson and Cooper 1976).

Anaerobic, sulfate-reducing bacteria (SRB) are the primary methylators of mercury in both lacustrine and estuarine sediments (Compeau and Bartha 1985; Gilmour and Henry 1991). The primary methylators of mercury in the water column have not been identified. SRB are common in sulfate-rich estuarine sediments (Hines et al. 1989) but are more limited in freshwater sediment with lower sulfate concentrations. A sulfate concentration of 200-500 µM in the water column is optimal for mercury methylation by SRB in sediment (Gilmour and Henry 1991). The activity of the methylating microbes is affected by environmental conditions (Jackson 1986) with nutrient availability and seasonality particularly important. The concentration of inorganic mercury in environmental media may not be a good indication of the concentration of methylmercury present due to the influence of environmental variables and biological activities. The importance of environmental factors in the production of methylmercury is as follows:

pH: Neutral or low pH conditions favor the production of monomethylmercury over dimethylmercury (Beijer and Jernelov 1979). An alkaline (high) pH favors the formation of dimethylmercury, which tends to escape into the atmosphere. Elevated tissue concentrations of methylmercury have been noted in numerous pristine lakes of the northern United States and Canada that receive acid rain and no point sources of mercury (Xun et al. 1987; Bloom et al. 1991). The mechanism(s) causing increased bioaccumulation in low pH lakes are not understood (Ramlal et al. 1985; Winfrey and Rudd 1990; Richardson and Currie 1996). The factors primarily responsible for net methylmercury production in lakes are, in decreasing order of importance, pH, dissolved organic carbon (DOC) concentration, and microbial respiration (Miskimmin et al. 1992). The importance of pH and sediment properties (Fe and Mn content) on methylation rates in saltwater environments has not been well studied.

Sulfide, Sulfate, and Other Ions: In the presence of sulfides, the mercuric ion (Hg[II]) becomes tightly bound to sulfide as insoluble mercuric sulfide and is not available for methylation. Sulfide activity may be the main factor influencing the availability of Hg[II] (Bjornberg et al. 1988) and the concentration of methylmercury in sediment (Craig and Moreton 1983). If pH is high or Eh is low, sulfide activity

will be high and mercury will be precipitated as insoluble mercuric sulfide. If the sulfide is oxidized to sulfate, the mercuric ion will become available for methylation. Both free sulfides and acid-volatile sulfides (AVS) appear to inhibit methylation (Gilmour and Capone 1987). The presence of other minerals may affect this relationship. Excess ferrous iron has been found to bind the sulfide and limit its Hg-binding effectiveness such that no difference in methylation rates are noted between sulfide-rich and sulfide-poor sediments (Rudd et al. 1983). Selenium similarly binds the Hg [II] ions and reduces their availability for methylation. The redox cycling of manganese in lakes may be more important than iron-scavenging of mercury (Bonzongo et al. 1996).

Addition of sulfate to anoxic lake sediment slurries or the overlying water column can increase methylmercury production by stimulating the SRB population (Gilmour et al. 1992). SRB can both methylate mercury and produce sulfide, which inhibits methylation: the kinetics of this are not understood. Mercury methylation was once viewed as a detoxification process in SRB, but it may ultimately serve some other function (Gilmour and Henry 1991).

Oxygen Conditions/Eh : Although methylation occurs under both aerobic (oxidizing) and anaerobic (reducing) conditions, methylation is greater under anaerobic conditions (Callister and Winfrey 1986; Weis et al. 1986; Regnell 1994). In addition, demethylation rates are lower under anaerobic conditions, so the net methylmercury production is higher in oxygen-depleted environments (Jackson 1987). Over 90 percent of methylmercury is formed biochemically in anaerobic sediment (Berman and Bartha 1986).

In anoxic lake bottoms containing hydrogen sulfide, mercury is bound to sediment as insoluble mercuric sulfide. If conditions become aerobic due to a decrease in the organic load or seasonal turnover, sulfide can be oxidized to sulfate, releasing the mercury in the ionic form Hg[II], which is available for methylation (Jernelov 1968).

Nutrients /Organic content :/DOC : Nutrients can enhance the rate of methylmercury production by stimulating the methylating bacteria. Decaying organic matter can enhance microbial activity and create low oxygen conditions, both of which cause higher methylation rates (Olson and Cooper 1974; Gilmour and Henry 1991). In freshwater areas with a high organic input, methylation rates

can become locally elevated if other environmental conditions do not inhibit methylation (i.e., high sulfide levels; Jackson 1986). When sulfide and sulfate concentrations are not limiting, organic matter may be the major factor controlling mercury methylation rates in estuarine sediments (Choi and Bartha 1994). Increased DOC levels may inhibit methylation due to the binding of free mercury ions (Jackson 1989; Winfrey and Rudd 1990) even though supplemental DOC increases microbial respiration (Miskimmin et al. 1992). In clear freshwater lakes, DOC and pH may interact such that less of the Hg [II] is bound by DOC at low pH, resulting in higher methylation rates. Acid rain may also limit the amount of DOC transported into a system because at lower pH, DOC solubility and mobility is reduced (De Haan 1992; Schindler et al. 1992).

Humic/Fulvic material: The geochemistry of Hg in lake and stream water may be dominated by humic material interactions (Mierle and Ingram 1991). Hg complexes with humic and fulvic substances and Hg retention and export from watersheds in Canada have been correlated with the export of humic substances. Hintelmann et al. (1995) assumed that the methylmercuric ion is bound to sulfidic binding sites of humic acid. At lower pHs, the amount of free unbound methylmercury ion was higher in their laboratory study of humic and fulvic acids. Acidification could potentially release bound methylmercury from humic acids into the aqueous phase where it would be readily bioavailable.

Salinity : There appears to be a negative correlation between the rate of methylmercury formation and salinity in estuarine sediments (Blum and Bartha 1980). The rate is lower in more saline environments because the bicarbonate component of seawater slows methylation of Hg [II] under both aerobic and anaerobic conditions (Compeau and Bartha 1983). The release of reactive Hg [II] and Hg [0] is slowed when chloride ions bind to mercury, thereby inhibiting methylmercury formation (Craig and Moreton 1985). Salinity also affects methylation due to the high pore-water sulfide concentrations as a result of rapid sulfate reduction in saline water compared to sulfate-limited freshwater environments (Gilmour et al. 1992). Along a salinity gradient in the lower Hudson River, methylation rates decreased downriver with increasing salinity and sediment sulfide concentrations (Gilmour and Capone 1987).

The percentage of total mercury that is methylmercury is higher in freshwater sediments (up to 37%) and water (up to 25% in aerobic water and 58% in anoxic bottom water) than in estuarine and marine water (<5%) and associated sediments (<5%) (Gilmour and Henry 1991). Dissolved reactive mercury (inorganic species) forms the majority of the total mercury in open oceans (Bloom and Crecelius 1983; Gill and Fitzgerald 1987).

Season: Biological productivity of methylating microbes is affected by seasonal changes in temperature, nutrient supply, oxygen supply, and hydrodynamics (changes in suspended sediment concentrations and flow rates). Methylmercury concentrations varied seasonally by an order of magnitude at most sites studied (Parks et al. 1989). Methylation may tend to increase during the summer months when biological productivity and temperature are high and decrease during winter months when biological productivity and temperature are low (Callister and Winfrey 1986; Jackson 1986; Weis et al. 1986; Korthals and Winfrey 1987; Parks et al. 1989; Kelly et al. 1995; Leermakers et al. 1995). Although the potential methylmercury production is greatest during the summer, actual production may not peak during this time (Kelly et al. 1995). In Onondaga Lake, New York, the mercury species in the water column varied temporally (Battelle 1987; Bloom and Effler 1990). Total mercury concentrations may also vary seasonally due to physical factors such as winter storms resuspending mercury-contaminated sediments (Gill and Bruland 1990).

Distribution in the Environment

The distribution and abundance of inorganic mercury and methylmercury in the environment may vary independently as they are controlled by different physicochemical processes. The concentration of total mercury (which is mainly inorganic) in the environment is generally not a good predictor of methylmercury concentration (Gilmour and Henry 1991; Kelly et al. 1995). Inorganic mercury has a high affinity for sediments; a significant portion of the total mercury in fresh water is in particulate form (Gill and Bruland 1990). Most of the mercury in estuaries was associated with particulate matter (Cossa and Noel 1987). The environmental distribution of inorganic mercury appears to be controlled by processes such as transport, sorting, and sedimentation as related to the hydrologic regime. Resuspension and resettling of sediments caused persistently high concentrations of mercury in the surface sediments of Lavaca Bay (Reigel 1990).

Total mercury concentrations in surface water may decrease as mercury bound to particulate matter settles or is transported downstream (Bonzongo et al. 1996). The distribution of biotically produced methylmercury initially depends on the microbial populations that methylate the mercury. Although more abundant in the sediment where it is formed, methylmercury forms a greater percentage of the total mercury in the overlying water column (Gilmour et al. 1992).

The distribution of both inorganic and methylmercury is also affected by the larger-scale physical characteristics of the environment such as type of system (river, lake, estuary, ocean) and its physical configuration, water circulation patterns, catchment type, sediment characteristics, rainfall, and the introduction of terrestrial sediments. The physical characteristics of the system influence the mechanisms of mercury distribution, availability, and cycling in varying degrees. Water flow regimes in particular subenvironments (mainstream versus backwater), characteristics such as the stratification cycle and amount of nutrients in lakes, or the configuration of the estuary (open circulation versus restricted), may have very different, distinct features that control both the persistence of mercury in the environment and how the different species of mercury behave and are distributed among sediment, water, and organisms.

In freshwater systems, Kelly et al. (1995) found a predictive, linear relationship between total and methylmercury concentrations in unfiltered water samples from some specific lake systems, but not from stream systems. Runoff from wetland catchments contributed more methylmercury to lake systems than did runoff from upland catchments (St. Louis et al. 1994). In estuarine systems, total dissolved mercury concentrations were found to be enhanced where salinity was less than 10 ppt, coinciding with the maximum turbidity zone (Cossa and Noel 1987; Cossa et al. 1988).

Terrestrial sediment influxes can also affect mercury availability. In the estuarine Ala Wai Canal in Hawaii, total mercury increased over two orders of magnitude in polychaetes and shrimp during the rainy season (Luoma 1977). Mercury bound to freshwater sediments and introduced into the estuary from urban runoff during rainfall was desorbed upon contact with saline water. The increased concentration of dissolved mercury temporarily increased total mercury in filter-feeding worms and shrimp. Total mercury concentrations in the water column and biota

decreased after the runoff stopped. In contrast, lower mercury concentrations were found in plankton in a freshwater lake after input of high concentrations of clean, fine-grained sediment. Sediments washed into the lake during rainfall bound the mercury, inhibiting uptake by plankton (Jackson 1988).

Sediment composition can also affect the way that mercury is distributed in the environment. Mercury concentrations in freshwater benthic organisms appeared to be determined by the sediment composition, such as the concentration of hydrated Fe and Mn oxides and carbon-rich humic matter in bottom sediment. The mercury appeared to be less available when it was bound by iron hydroxide (FeOOH), manganese hydroxide (MnOOH), and possibly by higher-molecular-weight humic substances (Jackson 1988).

In a freshwater river-lake system in Canada (Parks et al. 1989) methylmercury concentrations in surface water were highest 80 kilometers downstream from the most contaminated sediments (contaminated with inorganic mercury). Fish were contaminated as far as 270 km downstream from the inorganic mercury source, with the most contaminated fish found more than 100 km downstream of this source. Methylmercury concentrations in the water increased as inorganic mercury concentrations in the sediment decreased. The most highly contaminated sediments were located near a sewage outfall. The researchers surmised that the mercury in these sediments was bound to sulfide and thus not available for methylation. Further downstream the mercury became available for methylation probably due to a decrease in sediment sulfide levels. Even though concentrations of inorganic mercury in the sediment here were much lower than upstream concentrations, methylmercury production was much higher and biota were more contaminated.

There may be a similar situation in low-salinity water of Berrys Creek, New Jersey, where high inorganic sediment mercury concentrations were also found next to a sewage outfall (Weis et al. 1986). However, concentrations of mercury in fish inhabiting the area were not as high as expected. This was attributed to the presence of sulfide, which binds mercury and limits methylation. No downstream studies have been conducted to determine whether the mercury is more bioavailable further from the source of high sulfide concentrations.

An eight-ton cargo of elemental mercury located within the hold of the sunken *Empire Knight* in offshore marine waters did not contaminate invertebrates living outside the hold of the ship (Hoff et al. 1994). Only a small percentage of invertebrates sampled from within the hold had elevated concentrations of total mercury. However, it is not known whether the mercury was incorporated into the tissue of the organisms. The large source of elemental mercury in this environment was not bioavailable to organisms located away from the source.

BIOACCUMULATION OF MERCURY

Mercury bioaccumulates in aquatic plants, invertebrates, fish, and mammals. Concentrations increase (biomagnify) in higher-trophic-level organisms. Even though the different types of mercury have relatively low K_{OW} values (compared to organic compounds such as PCBs), they are readily accumulated. Inorganic mercury (excluding elemental) and methylmercury's strong reactivity with intracellular ligands is thought to be responsible for their high degree of accumulation. Uptake and accumulation of mercury are affected by the type of mercury present, with neutral mercury species (e.g., $HgCl_2^0$ and CH_3HgCl^0) absorbed more efficiently than charged mercury species (e.g., $HgCl^-$ and CH_3Hg^+ ; Mason et al. 1996).

Despite the fact that the neutral inorganic and organic complexes have similar lipid solubilities, methylmercury is selectively accumulated (due to a higher transfer efficiency and lower rate of elimination), resulting in biomagnification in higher trophic levels (Mason et al. 1995b). Inorganic mercury species are not biomagnified (Surma-Aho and Paasivirta 1986; Riisgård and Hansen 1990; Hill et al. 1996).

Environmental factors that enhance mercury methylation result in greater bioavailability and accumulation of methylmercury. Environmental variables also influence the bioavailability and accumulation of inorganic mercury. Although concentrations of mercury in the environment may correlate with concentrations in resident plants and biota, correlation is often difficult. Correlating total mercury in sediment with total mercury in upper-trophic-level organisms is complicated by high methylmercury concentrations in high-trophic-level organisms relative to low methylmercury concentrations in the environment.

Tissue concentrations of mercury are often positively correlated with organism length, weight, and/or age. Diet has a significant role in the overall body burden of mercury, both between and within species. Differences in total mercury concentrations between species reflect diet differences due to trophic position; within-species differences are related to dietary requirements of various developmental stages.

The Effect of the Form of Mercury on Bioaccumulation

Both inorganic and methylmercury are taken up directly from water and food (or ingested sediment). However, methylmercury is more efficiently accumulated than inorganic mercury for most aquatic organisms (Fowler et al. 1978; Julshamn et al. 1982; Riisgård and Hansen 1990; Mason et al. 1995b). The uptake and depuration of mercury depends on the form of mercury, source of mercury (water or food), and the type of receptor tissue, resulting in different patterns of accumulation. Methylmercury is readily transferred across biological membranes. Within the organism, methylmercury is strongly bound to sulfhydryl groups in proteins of tissues such as muscle, and is much slower to depurate than inorganic mercury. Thus, methylmercury has a much greater potential for bioaccumulation and a longer half-life in organisms than inorganic mercury.

Fish

The accumulation of mercury from water occurs via the gill membranes. Gills take up aqueous methylmercury more readily than inorganic mercury (Huckabee et al. 1979; Boudou et al. 1991). Methylmercury is eventually transferred from the gills to muscle and other tissues where it is retained for long periods of time (Julshamn et al. 1982; Riisgård and Hansen 1990).

Inorganic mercury taken up with food initially accumulates in the tissues of the posterior intestine of fish (Boudou et al. 1991). Inorganic mercury is not easily transferred through this organ to other parts of the body. After 15 days, 80% had depurated from the fish intestine. Liver and kidney in fish tend to have higher percentages of inorganic mercury than muscle tissue, although percentages vary by organ and species (Windom and Kendall 1979; Riisgård and Hansen 1990).

Methylmercury ingested in food is efficiently transferred from the intestine to other organs (Boudou et al. 1991). Methylmercury has been reported to constitute from 70 to 95% of the total mercury in skeletal muscle in fish (Huckabee et al. 1979; EPA 1985; Riisgård and Famme 1988; Greib et al. 1990; Spry and Wiener 1991).

Methylmercury accounted for almost all (99%) of the mercury in muscle tissue in a wide variety of both freshwater and saltwater fish found in waters not highly contaminated by other organomercurial species (Bloom 1992).

The ratio of liver to muscle total mercury concentration usually fluctuates around one and can reflect the exposure history of the organisms. For example, the liver:muscle ratio may be less than one in chronically exposed fish, while a recent exposure to mercury may result in a ratio greater than one (Riisgård and Hansen 1990).

McKim et al. (1976) reported that mercury could be transferred from adult to offspring in brook trout. Exposure of the parent population to aqueous methylmercury concentrations of 0.03 to 2.93 $\mu\text{g}/\text{l}$ in the laboratory resulted in mercury concentrations as high as 2 mg/kg in their embryos. Total mercury concentrations in eggs of several species of adult fish from Swedish lakes were much lower than concentrations in other tissues; therefore, spawning did not lower their total mercury body burden (Lindqvist 1991).

The main depuration pathway is through the kidney and liver in fish. Half-lives for methylmercury in fish range from one to three or more years (McKim et al. 1976; Pentreath 1976a, b; Riisgård and Famme 1986; Riisgård and Hansen 1990), while estimates of half-lives for inorganic mercury are much lower, ranging from approximately five days to five months (Pentreath 1976a, b; Huckabee et al. 1979).

Invertebrates

Invertebrates accumulate and partition inorganic and methylmercury in tissues similar to the trends exhibited by fish (Fowler 1978; Riisgård and Famme 1986; Saouter et al. 1991; Saouter et al. 1993). However, invertebrates generally contain a lower percentage of methylmercury than fish or mammals (Lasorsa and Allen-Gil 1995), with highly variable concentrations. This wide variation of mercury content in invertebrates is most likely a function of different feeding strategies (and trophic

levels) and different environmental exposures. Reported percentages of methylmercury compared to total mercury concentrations are less than 1% for the polychaete *Nereis succinea* (Luoma 1977); 10% in copepods, mussels and shrimp (Horvat 1991); 10-100% in the cockle (Møhlenberg and Riisgård 1988); 16% in urchin gonads (Eganhouse and Young 1978); 30-90% in lake zooplankton (Lindqvist 1991); 87% in crab muscle (Eganhouse and Young 1978); and 100% in red rock crab, Dungeness crab, and spot shrimp (Bloom 1992). Becker and Bigham (1995) found an increasing percentage of methylmercury compared to total mercury in higher trophic levels in the Onondaga lake food web. Lake water contained 5% of total mercury as methylmercury; phytoplankton 24%; benthic macroinvertebrates 26%; zooplankton 40%; and fish fillets 96%.

Viscera in mussels contained the highest tissue concentration of total mercury (Fowler et al. 1978). The total mercury concentration was highest in the midgut and muscle tissue in crab (Bjerregaard and Christensen 1993) and in the viscera in shrimp (Fowler et al. 1978). The shrimp molts had the lowest mercury content; therefore, molting is not considered an important depuration pathway in crustaceans (Fowler et al. 1978).

Half-lives for total mercury in salt-water mussels ranged from two months to one year (Riisgård et al. 1985). Inorganic mercury was eliminated more rapidly than methylmercury in mussels and shrimp (Fowler et al. 1978).

Marine mammals

Marine mammals have some of the highest tissue mercury concentrations of all marine organisms investigated (Andre et al. 1991a); however, concentrations are highly variable both within and among species. These variations have been attributed to collection locations (Wren 1986), concentrations in prey items (Szefer et al. 1993), and organism age (Julshamn et al. 1987; Thompson 1990). For example, species that feed primarily on benthic invertebrates, such as walrus and baleen whales, tend to have relatively low mercury concentrations. In contrast, fish-eating species, such as porpoises and seals, exhibit relatively high mercury concentrations (Born et al. 1981).

In contrast to fish, adult marine mammals have a much higher percentage of total mercury as inorganic mercury, although the concentration of methylmercury may also be elevated. Less than 10% of the total mercury content is methylmercury (Eisler 1987). Juveniles tend to have higher percentages of methylmercury.

The liver generally exhibits the highest total and methylmercury concentration (Holden 1978; Wagemann et al. 1983; Julshamn et al. 1987; Thompson 1990; Andre et al. 1991a), followed by kidney and muscle tissues (Szefer et al. 1993). Julshamn et al. (1987) measured the highest concentrations of methyl- and total mercury (13 and 150 mg/kg) in pilot whale livers (*Globicephalus meleanus*) compared to total and methylmercury concentrations in muscle (2.8 mg/kg total mercury; 1.7 mg/kg methylmercury) and kidney (15.3 mg/kg total mercury; 5.1 mg/kg methylmercury). Andersen et al. (1987) also measured the highest methylmercury concentrations in pilot whale liver (20 mg/kg; 14% of total mercury). The fraction that is methylated, however, is usually lower in the liver compared to muscle and kidney. The methylated fraction of total mercury ranged from 1% to 36% in seal liver (Holden 1978); 30% in older specimens to 100% in young specimens in the muscle of harbor porpoise (Joiris et al. 1991); and 24% to 86% in the muscle of pilot whales (*Globicephalus meleanus*; Julshamn et al. 1987). The liver:muscle ratio for methylmercury concentration in harbor porpoises was approximately one, while the ratio for liver:muscle for total mercury concentration was two. In some of the harbor porpoises and some other species (sperm whale, common dolphin, and adult bottle-nose dolphin) the liver:muscle ratio for total mercury ranged up to 20 (Joiris et al. 1991) while the liver:muscle ratio for methylmercury was still one. Schintu et al. (1992) observed an age-related change in the percentage of methylmercury compared to total mercury in pilot whale livers. The liver of three- to seven-year old pilot whales (with a relatively low total mercury body burden) contained 30% to 60% organic mercury, compared to 3% to 17% organic mercury in livers of 30- to 40-year old pilot whales (with a relatively high total mercury load). Porpoises exhibited a similar trend. Juveniles had a higher percentage of methylmercury in liver (100%), while the percentage of methylmercury in adult specimens decreased to 2 to 3% of the total mercury (Joiris et al. 1991).

Although mercury in the diet of many marine mammal species is predominantly methylmercury, it has been proposed that the mammals are able to mineralize methylmercury into the more harmless inorganic form, which then accumulates in the liver of adult specimens (Holden 1978; Joiris et al. 1991). The estimated half-life

of total mercury in pinnipeds and dolphins is about 1.4 and 2.7 years, respectively (Eisler 1987; Andre et al. 1991b).

Plants

Vascular plants accumulate both inorganic and methylmercury from sediment and water in root, stem, and leaf sections (Alberts et al. 1990; Boudou et al. 1991). The rooted macrophyte *Elodea densa* accumulated different concentrations of methylmercury versus inorganic mercury from sediment (uptake was 40 times higher for methylmercury). Tissue concentrations were similar throughout the plant when the mercury source was water (ratio of 1.5 methylmercury to inorganic; Boudou et al. 1991).

Chloride concentration and pH influenced uptake of inorganic mercury by a marine diatom. Rates were low in seawater and low-chloride freshwater with neutral pH (Mason et al. 1996). Methylmercury uptake rates were high in high-chloride waters and were not influenced by pH. The uptake rate of methylmercury only became limited when very low chloride concentrations decreased the concentration of CH_3HgCl . Elemental and dimethylmercury were not significantly accumulated.

Exposure Pathways

Aquatic organisms can accumulate mercury from water (including pore water) and food sources (including sediment). Quantity accumulated is a function of the exposure pathway and the physical and environmental factors such as temperature, pH, salinity, total organic carbon, and sulfides. If conditions are favorable for methylation, organisms can accumulate high concentrations of mercury even with low concentrations in the water and sediment.

Water

Phytoplankton, invertebrates, fish (including eggs and larvae), and mammals take up inorganic and organic mercury from the water column (McKim et al. 1976; Pentreath 1976a; 1976b). In phytoplankton, algae, and microorganisms, mercury uptake is primarily a passive process that occurs by adsorption to the cell surface either through interaction with functional groups in the cell wall or through sorptive properties associated with the extracellular matrices (Darnell et al. 1986; Gadd 1988). Passive diffusion of lipid-soluble species (uncharged chloride

complexes) is responsible for mercury uptake in a marine diatom (Mason et al. 1996). Uptake in phytoplankton and aquatic plants has been correlated with the concentration of mercury in the water (Windom and Kendall 1979; Lenka et al. 1990). Water is an important exposure pathway for mercury uptake by lower organisms and thus into the food web (Francesconi and Lenanton 1992). Dissolved mercury concentrations in water are typically very low; the major increase in mercury concentrations occurs between water and phytoplankton of about a factor of 10^5 to 10^6 (Mason et al. 1995b). In contrast to microorganisms, uptake is primarily an active process for fish and invertebrates, and is related to respiration rate and metabolic rate (Rodgers and Beamish 1981). Uptake of methylmercuric chloride in water by different tissues of brook trout was found to be directly related to the water concentration of the mercury (McKim et al. 1976).

BCFs are the concentration of mercury in tissue divided by the concentration in the exposure water. They have been calculated from laboratory experiments for many species of aquatic organisms to estimate uptake from water. However, BCFs have limited use for several reasons. First, BCFs reported in the literature most likely underestimate actual values because laboratory studies were done before the use of trace-metal free protocols and used higher water concentrations than found in the field (Zillioux et al. 1993). More recent BCF calculations for mercury have yielded values one to two orders of magnitude higher than previous estimates. Second, BCFs only reflect uptake of a contaminant from the water. Higher trophic species accumulate mercury primarily through the food web. Reported BCFs for mercury vary considerably due to differences between species, exposure concentration, and duration. Further, BCFs for the same species may be several orders of magnitude higher for methylmercury than for inorganic mercury.

Brook trout exposed to varying concentrations of methylmercury for 28 to 38 weeks had bioconcentration factors ranging from 69,000 to 630,000 (McKim et al. 1976). The wide range of BCFs reported for brook trout are related to the tissue analyzed. Bioconcentration factors for muscle tissue in brook trout were higher at lower water concentrations of mercury. BCFs for other tissues remained the same when water methylmercury concentrations were varied.

Bioconcentration factors for inorganic mercury (mercuric chloride) in saltwater species were 129 for adult lobster (*Homarus americanus*), 1,000 for mussels, and 10,000 for oysters (*Crassostrea virginica*; Kopfler 1974; Roesijadi et al. 1981). Bioaccumulation factors (BAFs) were calculated from field studies for yearling yellow perch from five freshwater lakes. These factors ranged from 10^6 - 10^7 for methylmercury and more than 10^4 for other mercury species (Bloom 1992).

Sediment

Sediment is an important exposure pathway for all forms of mercury to aquatic organisms. High concentrations of organic substances and reduced sulfur that complex free Hg[II] ions in sediment can reduce the availability of mercury to biota (Luoma 1977; Rubinstein et al. 1983). Correlating mercury concentrations in sediment with concentrations in biota may be difficult, particularly for higher-trophic-level species.

The bioavailability of total mercury to benthic invertebrates was reported to be inversely correlated to the organic content of the sediment (Langston 1982, 1986). Normalizing sediment mercury concentrations to percent organic matter improved the correlation between total mercury concentrations in sediment and invertebrate species (including gastropods, polychaetes, and deposit- and suspension-feeding bivalves) in a marine environment (Bryan and Langston 1992). Good sediment-tissue correlations for mercury have been found in amphipods from a freshwater lake (Becker et al. 1993). Breteler et al. (1981) studied mercury uptake by plants and invertebrates from several types of sediments in salt marsh environments. Concentrations of total mercury in mussels, fiddler crabs, and *Spartina alterniflora* increased as organic matter in sediments decreased.

Many investigators report no correlation between sediment and tissue concentrations of mercury for higher-trophic-level species (Nishimura and Kumagai 1983; Jackson 1988; Rada et al. 1989b; Lindqvist 1991; Duckerschein et al. 1992). Organic carbon normalization of sediment concentrations did not improve the correlations for pike, a high trophic level species (Lindqvist 1991). The difficulty in correlating mercury in sediment with mercury in organisms reflects the complexity of variables that affect both the methylation of mercury in surface sediments and the transfer of mercury between trophic levels. Since methylation occurs primarily

in surface sediments, the physical factors that affect the rate of methylation (and demethylation) also affect the availability of mercury for uptake by organisms. Sediment total-mercury concentrations alone may not provide information on the exposure potential of resident organisms.

Food web

Though sediment may be the ultimate source of mercury for many higher trophic species, the food web is the primary pathway to most organisms (Lindqvist 1991; Bryan and Langston 1992). Most of the differentiation between inorganic and methylmercury accumulation occurs during trophic transfer (Mason et al. 1995b) because of the differences in assimilation of the different mercury forms and how efficiently the different forms are transferred to predators.

Mason et al. (1995b) detected an assimilation efficiency four times greater for methylmercury compared to inorganic mercury from phytoplankton to zooplankton, and ten times greater between phytoplankton and planktivorous fish. The transfer efficiency of methylmercury over inorganic mercury in zooplankton was attributed to mercury partitioning in the algal cell. Methylmercury accumulated in the algal cytoplasm, which zooplankton digest, with 62% of the methylmercury transferred, while inorganic mercury was primarily bound to thiols in the algal cell membrane. Therefore, a smaller percentage (15%) of inorganic mercury was transferred to zooplankton.

As methylmercury increases in prey items, the transfer efficiency also increases (Windom and Kendall 1979). Since methylmercury concentrations are highest in fish, piscivorous fish will be exposed to higher concentrations of methylmercury than fish that feed on invertebrates. For example, walleye accumulated mercury at a faster rate and at higher concentrations than pike from the same freshwater lake (Mathers and Johansen 1985). A high proportion of the diet of walleye was smelt, the most contaminated prey item, whereas pike ate only a small proportion of this prey item. Dietary changes during life history development, or due to season or habitat differences can change exposure. Dietary shifts in prey items of similar trophic levels but from different habitats, or dietary shifts due to a different size structure of prey, can also affect the mercury concentrations in top-level predators (Lindqvist 1991).

The relative importance of dietary versus aqueous mercury uptake pathways is unclear. Probably less than 10% of the mercury in fish tissue residues is obtained by direct (gill) uptake from water (Francesconi and Lenanton 1992; Spry and Wiener 1991). Methylmercury concentrations used in laboratory studies of aqueous uptake are 1,000 to 10,000 times the ambient concentration of methylmercury in natural water (Spry and Wiener 1991), thereby overestimating the significance of direct aqueous uptake. The proportion of mercury taken up from dietary sources versus water in invertebrates has not been estimated. Suspension-feeding bivalves may principally accumulate mercury by consuming algal cells (Riisgård and Hansen 1990).

Although mercury correlations are complicated by the importance of the food-chain exposure pathway, mercury concentrations in predators and prey have been correlated (e.g., Allard and Stokes 1989; Lindqvist 1991; Spry and Wiener 1991). For example, mercury concentrations in smallmouth bass from Ontario lakes were directly correlated with mercury in crayfish, which comprised 60% of their diet.

Detritus can be a very important source of mercury, particularly in estuarine habitats. Organic detritus from *Spartina alterniflora* may contain 30 times more mercury than plankton. Organisms in detritus-based food webs are thus exposed to higher mercury concentrations than are animals feeding on plankton (Lindberg and Harriss 1974). Mercury associated with humic matter in lakes is fed upon by bacteria and zooplankton, which incorporate mercury into the detrital food web (Lindqvist 1991).

Mercury in the fecal matter of marine mammals can also be a significant source to other aquatic organisms near breeding colonies or haul-out areas (Eisler 1987).

Biological Factors Affecting Accumulation of Mercury

The primary biological factors governing the accumulation of mercury include age, weight, and diet. Differences in accumulation between the sexes have been attributed to differences in diet.

Fish

Numerous field studies have shown that the concentration of total mercury in fish positively correlates with length, age, and weight (Hall et al. 1976a, b; Huckabee et al. 1979; Rada et al. 1986; Møhlenberg and Riisgård 1988; Greib et al. 1990; Leah et al. 1992). However, total mercury concentrations may not always correlate with size due to differences associated with diet, residence time in a contaminated habitat, and type of mercury (Francesconi and Lenanton 1992). The percentage of methylmercury increases with age in both fish and invertebrates (Møhlenberg and Riisgård 1988; Riisgård and Hansen 1990).

In some species of fish and invertebrates, sex differences in mercury tissue concentrations have been reported. For example, total mercury concentrations in the muscle tissue of freshwater sunfish were greater in females than males at ages 2 to 3 (Nicoletto and Hendricks 1987). This may be due to increased food demands for females related to reproduction. In contrast, there was little relationship between sex and bioaccumulation of mercury in three species of fish (roach, perch, and pike) collected from Swedish lakes (Lindqvist 1991).

Bloom (1992) did not find a relationship between lipid content and methylmercury concentrations in a variety of fresh- and saltwater fish.

Invertebrates

Cockles (*Cardium edule*) from a polluted estuary were found to have a positive linear correlation between their age and the percentage of organic mercury in their tissues (Møhlenberg and Riisgård 1988). Organic mercury comprised 30% of the total mercury in two-year old cockles; 60% in three-year olds; and 90% in four-year olds. This relationship was attributed to the rapid loss of inorganic mercury and continued uptake of organic mercury over time. However, the correlation was not as strong when weight was used instead of age due to variations in growth rate at different locations. Total mercury concentrations in mussels were found to be higher in 27 mm than 31-mm sized individuals (Riisgård and Hansen 1990). This difference was perhaps due to a decrease in both weight-specific filtration rate and surface area-to-volume ratio in larger mussels (Fowler et al. 1978; Riisgård et al. 1985).

Total mercury concentrations may (Allard and Stokes 1989) or may not (Rada et al. 1986) correlate with weight or age in crustaceans.

Concentrations of mercury in male and female emergent mayflies (*Hexagenia bilineata*) in the upper Mississippi River differed. The authors recommend sampling male and female mayflies separately (Dukerschein et al. 1992).

Marine Mammals

Mercury concentrations (both organic and inorganic) are positively correlated with body length in marine mammals (Arima and Nagakura 1979; Wagemann et al. 1983; Joiris et al. 1991). Hansen et al. (1990) found a highly significant correlation between age and tissue content of mercury in whales from West Greenland. This correlation has been used to separate immature specimens from adults. Joiris et al. (1991) found that the concentration of methylmercury in muscle and liver tissue in harbor porpoises did not increase with increasing length as strongly as did total mercury.

Leonzio et al. (1992) suggest that the elevated concentrations of inorganic mercury measured in mammals, as compared to fish, may be related to differences in respiratory systems. In contrast to fish, where the gills allow contaminants to be lost to the environment because blood flow has contact with the water, the mammalian respiratory system does not have a similar exchange. Mammals have developed different defense mechanisms. For example, selenium combines with mercury to form the non-toxic compound tiemannite that is stored within cells. The processes of intracellular storage tend to increase concentrations of the metal in certain organs while reducing the toxicity. In marine mammals, intracellular storage of mercury occurs as complexes of both selenium and metallothioneins (MTs).

Other Factors Affecting Accumulation

Temperature and season influence the availability and accumulation of mercury in addition to the factors already discussed. Changes in temperature can affect mercury concentrations in organisms either directly by affecting metabolic rate and thereby exposure, or indirectly by influencing the methylation of mercury and therefore enhancing availability. Rates of methyl- or inorganic mercury uptake

increase with increasing aqueous concentrations and/or increasing temperature in the water for some species (e.g., phytoplankton, gastropods, fish; Windom and Kendall 1979; Rodgers and Beamish 1981; Tessier et al. 1994). A rise in temperature (and a corresponding rise in respiratory volume) can increase the rate of uptake via the gills (EPA 1985).

Total concentrations of mercury in killifish from an estuarine wetland were five times higher in spring and summer than in other seasons (Weis et al. 1986), presumably due to higher methylation rates in summer. Zooplankton mercury concentrations peaked in June in Swedish lakes and fish tissue levels varied by a factor of two, reaching a maximum in spring (Lindqvist 1991). Mercury content of mussels from the Gulf of St. Lawrence estuary varied seasonally by a factor of two (Cossa and Rondeau 1985).

The relationship of pH, conductivity, and salinity to mercury accumulation is not well understood. Elevated mercury concentrations have frequently been found in piscivorous fish in poorly buffered (alkalinity < 55 µeq/l and calcium < 2 mg/l), low-pH lakes (pH 6.0-6.5) in areas removed from industrial inputs of mercury (Rada et al. 1989a; Winfrey and Rudd 1990; Spry and Wiener 1991). Total mercury concentrations in yellow perch were inversely correlated with pH in ten Wisconsin lakes (Cope et al. 1990). Mercury concentrations in zooplankton in Swedish lakes were correlated with pH but the relative importance of this correlation changed over time (Lindqvist 1991).

In freshwater lakes removed from direct sources of mercury, conductivity explained 54% of the variability in mercury concentrations in crayfish (Allard and Stokes 1989). Conductivity was also highly correlated with calcium, magnesium, alkalinity, pH, and sodium. This correlation suggests that the buffering capacity of the lake was an important influence on crayfish accumulation of mercury. Low calcium ion concentrations enhanced the efficiency of methylmercury uptake across the gills of rainbow trout (Rogers and Beamish 1983).

TOXICITY OF MERCURY

The toxicity of mercury to aquatic organisms is affected by both abiotic and biotic factors including the form of mercury (inorganic versus organic), environmental conditions (e.g., temperature, salinity, and pH), the sensitivity of individual species and life history stages, and the tolerance of individual organisms. Toxicological effects include neurological damage, reproductive impairment, growth inhibition, developmental abnormalities, and altered behavioral responses. Wiener and Spry (1996) concluded that neurotoxicity seems to be the most probable chronic response of wild adult fishes to methylmercury exposure, based on observed effects such as uncoordination, inability to feed, diminished responsiveness, abnormal movements, lethargy, and brain lesions. In laboratory studies, reproductive endpoints are generally more sensitive than growth or survival, with embryos and the early developmental stages being the most sensitive (Hansen 1989). Impaired reproduction in sensitive aquatic organisms has been shown to occur at aqueous concentrations of mercury between 0.03 and 1.6 $\mu\text{g/l}$ (Eisler 1987). Long-term mercury exposure to adult fish also has been shown to result in retarded growth of offspring (Snarski and Olson 1982) and teratogenic effects (Weis et al. 1981). Chronic exposure to low concentrations of mercury may result in populations that become tolerant to the toxic effects of mercury contamination (Weis and Weis 1989).

The toxic concentration of mercury compounds can vary by an order of magnitude or more depending on the exposure condition. For example, toxicity is greater at elevated temperatures (Armstrong 1979), at lower oxygen content (Sloof et al. 1991), and at reduced salinities in marine environments (McKenney and Costlow 1981). Site-specific factors (such as TOC) affect the bioavailability and toxicity of mercury-contaminated sediment (Langston 1990). Even though correlations exist between toxicological observations and sediment pollution gradients, Langston (1990) recommends collecting site-specific data because biological responses can not always be satisfactorily predicted from chemical data or modeling results.

The sensitivity of aquatic organisms to either inorganic or methylmercury varies considerably between species – more than the difference in sensitivity of a particular species to various mercury compounds (EPA 1985). Methylmercury is more acutely toxic to aquatic organisms than inorganic mercury, but the range

among different species in sensitivity to either compound is quite large. For example, the concentration of inorganic mercury inducing acute toxicity was observed to range over almost three orders of magnitude from 0.1 µg/l to more than 200 µg/l when results from tests with different species were compared (Eisler 1987). Tests on the same freshwater species with both inorganic and methylmercury showed that methylmercury was more than 30 times more acutely toxic than inorganic mercury (EPA 1985).

The general mechanism of action for toxic effects for inorganic mercury which has the form Hg (II), the divalent mercury cation, is believed to be the high affinity for thiol or sulfhydryl groups of proteins (Clarkson 1972; Hughes 1957, Passow et al. 1961) resulting in altered protein production or synthesis (Syversen 1977). Methylmercury is lipid soluble, allowing rapid penetration of the blood-brain barrier (Feltier et al. 1972, Giblin and Massaro 1973; McKim et al. 1976; Olson et al. 1978; Beijer and Jernelov 1979). Injury to the central nervous system results from accumulation of methylmercury in the cerebellum and cerebral cortex where it binds tightly to sulfhydryl groups resulting in pathological changes (Sastry and Sharma 1980). Inside the cell, methylmercury inhibits protein synthesis/RNA synthesis (Yoshino et al. 1966; Chang et al. 1972).

Zillioux et al. (1993) suggest that, prior to the mid-1980s, few data are available on the biological effects of mercury at environmental concentrations because laboratory studies used exposure concentrations that were much higher than actual concentrations in the field. This was in part due to contamination during sample collection and analysis. Improvements in trace-metal-free clean protocols during sample collection, handling, and processing as well as lower analytical detection limits have resulted in lower environmental concentrations of mercury and lower concentrations reported to elicit adverse effects. Although pre-1980 data are useful in identifying modes of effect and relative toxicity of the various mercury compounds, these data should be used with caution.

The following sections review available literature on the toxicity of mercury in water, toxicity in sediment, and toxicity associated with mercury in tissues. A wide range of toxic concentrations have been reported.

Toxicity of Mercury in Water

Nearly all of the studies evaluating the toxicity of mercury compounds where the route of exposure is through water have been conducted under laboratory conditions. Due to the nature of laboratory studies and differences in experimental design and technique, a wide range of toxic concentrations have been reported for a given species (Table 1). For example, toxicity tests using flow-through systems generally show higher toxicity at lower concentrations than static-renewal systems using the same (nominal) concentrations and the same species. This difference is probably due to loss of mercury from the test container in the static-renewal tests (Birge et al. 1979; Biesinger et al. 1982; WHO 1989).

Fish

Fish tend to be more sensitive to sublethal effects from chronic exposure to both inorganic and organic mercury than invertebrates, but fish are less sensitive to acute effects (EPA 1985; Hansen 1989). The early life stages of fish are generally the most sensitive to mercury. Birge et al. (1979) conducted several tests designed to evaluate embryo survival, hatching success, teratogenic effects, and the effects of mercury on six species of freshwater fish. The sensitivity of the embryo-larval stage for various species was correlated with the length of time for eggs to develop and hatch and the duration of exposure. Trout eggs treated in a flow-through system experienced approximately 40% mortality after a five-day exposure and 100% mortality after an eight-day exposure to an average mercury concentration of 0.12 µg/l.

Birge et al. (1979) also evaluated the long-term effects of mercury exposure on fish reproduction by conducting chronic bioassays with rainbow trout. Their results suggest that exposure of adult fish to mercury can have significant adverse effects on their offspring, with the effects enhanced if the embryos are also reared in a mercury-contaminated environment. Their data show a dose-dependent response in both bioaccumulation of mercury in gonadal tissues and toxic effects on embryos. Short-term exposures of embryos to high concentrations of mercury can also elicit significant adverse effects (Sharp and Neff 1980) and such exposures should be taken into account in the evaluation of potential long-term impacts to receiving environments. Although time-integrated concentrations may be within accepted guidelines, a short-term exposure to an elevated mercury concentration

Table 1. Toxicity resulting from exposure to mercury in water.

Fish Species	Hg ($\mu\text{g/l}$)	Hg Form	Exposure	Effect(s)	Reference
Brachydanio rerio Zebrafish (adults & fertilized eggs)	0.1, 5, 10, & 20	phenylmercuric acetate	25 days	83% spawning inhibition at 5 $\mu\text{g/l}$; 99% inhibition at 20 $\mu\text{g/l}$; significant decrease in hatching frequency at 0.2 and 1 $\mu\text{g/l}$.	Kihlstrom et al. 1971
Brachydanio rerio Zebrafish (fertilized eggs)	10, 20 & 50	phenylmercuric acetate	~ 6 days	increased hatching success at 10 $\mu\text{g/l}$; increased time to hatch and reduced hatching success at 20 $\mu\text{g/l}$; no hatch at 50 $\mu\text{g/l}$	Kihlstrom and Hulth 1972
Poecilia reticulata Medaka guppies	1.8	methylmercury	3 months	impaired spermatogenesis	Wester 1991
Oryzias latipes Medaka (fertilized eggs)	15	mercuric chloride	16 days	80% egg mortality; cardio-vascular abnormalities in eggs (hemorrhaging, blood vessel deterioration, loss of circulating blood cells and blood vessels), teratogenic effects in fry (i.e., non- functioning caudal fins, skeletal defects, spinal curvature)	Heisinger and Green 1975
Fundulus heteroclitus Killifish (fertilized eggs)	20 & 30 0, 4, 10, 20, 30, 40, 60, 80 & 100	mercuric chloride	32 days	100 % egg mortality decreased hatching at $[\text{Hg}] > 10 \mu\text{g/l}$; lateral curvature of the spine in larvae at $[\text{Hg}] 30 \mu\text{g/l}$.	Sharp and Neff 1980
Fundulus heteroclitus Killifish (fertilized eggs)	30 & 40	methylmercuric chloride	up to 48 days	reduced axis formation, development of cycloopia, defective cardiovascular system, skeletal malformations	Weis and Weis 1977
Pimephales promelas Fathead Minnow	0.26 1.02 2.01	mercuric chloride	41 days	decreased growth in females and reduction in number of spawning pairs cessation of spawning severe stunting and scoliosis	Snarski and Olson 1982
Oncorhynchus mykiss Rainbow trout (fertilized eggs)	0.12	mercuric chloride	5 & 8 days	40% mortality in fertilized eggs; 100% mortality after 8 days	Birge et al. 1979

Table 1. continued

Molluscs/Bivalves Species	Hg ($\mu\text{g/l}$)	Hg Form	Exposure	Effect(s)	Reference
Crepidula fornicata Limpet	>0.25 0.42	mercuric chloride	16 weeks	growth impairment, diminished condition suppressed fecundity, reductions in larval settlement	Thain 1984
Mercenaria mercenaria Clams	1.0 15	mercuric chloride	8 & 10 days	significant growth reductions reduced growth	Calabrese et al. 1977
Crassostrea virginica Oysters	12			reduced growth	
Mytilus edulis Mussels	0.3 >1.6	mercuric chloride	10-22 days >3 days	reduced growth cessation of growth	Stromgren 1982
Echinoderm Species					
Anthocardia crassispina Sea urchin	10	mercuric chloride	Not available	fertilization and development interference	Kobayashi 1984
Crustacean Species					
Callinectes sapidus Blue crab (megalopae through 2nd crab stage)	10 to 20	mercuric chloride	22 days	reduced survival in megalopae	McKenney and Costlow 1981
Mysidopsis bahia Mysids	1.6 2.5	mercuric chloride	full life cycle	delayed sexual maturation, delayed brood release, decreased brood production increased brood development time; aborted developing juveniles	Gentile et al. 1983

could result in inhibition of hatching, teratogenic development, and possible population effects.

Low concentrations of mercury in freshwater reportedly result in olfactory and chemoreceptor impairment in salmonids and other fish, which may interfere with normal migratory behavior (Hara et al. 1976; Rehnberg and Schreck 1986). For example, Hara et al. (1976) reported that rainbow trout exposed to inorganic mercury concentrations as low as 0.1 mg/l for two hours showed reduced olfactory responses. Further physiological and behavioral studies by Rehnberg and Schreck (1986) showed that mercury exposure reduced the ability of coho salmon to detect natural odors and disrupted simple upstream movement in laboratory experiments.

Weis and Weis (1989) suggest that prior exposure to mercury may produce populations that are more tolerant to the toxic effects of mercury contamination. Differences in tolerance to the effects of methylmercury were observed between organisms from mercury-contaminated and clean environments. Eggs collected from killifish in a contaminated area were mostly resistant to the teratogenic effects of methylmercury, while eggs of fish from the clean area showed a range of sensitivity. The susceptible eggs from the clean area also accumulated higher levels of mercury than did the eggs from the contaminated area (Weis et al. 1981; 1982). Offspring from fish previously exposed to mercury contamination were more tolerant to environmental mercury concentrations than offspring from clean environments (Weis and Weis 1984).

The situation is complicated by the fact that some fish that build up a tolerance to low concentrations of mercury can also detoxify the free metal within cells via the production of metallothioneins (MTs) and other metal-binding proteins. Brown et al. (1983) proposed that toxic effects occur as the binding capacity of MT becomes saturated, due to the interaction of excess free metal in the cell with the enzyme pool.

Invertebrates

Calabrese et al. (1977) suggest that marine bivalves embryos are more sensitive than the larvae in their susceptibility to mercury. They further indicated that growth of fully-developed larvae may be retarded at concentrations too low to elicit

significant mortality, thus prolonging the pelagic stages and increasing the risk of predation, disease, and dispersion. Several endpoints have been used to measure the effect of mercury exposure on bivalves, including biomarkers. The prophyrin precursor δ -aminolevulinic acid (ALA) may be useful as a biomarker of mercury exposure in bivalves (Brock 1993).

The effects of salinity on the toxicity of mercury have been demonstrated in a study conducted with the megalopae of the blue crab, *Callinectes sapidus* (McKenney and Costlow 1981). Their results indicated that as salinity was reduced below 20 parts-per-thousand, less mercury was required to produce equivalent toxicity among megalopae. This is significant for blue crab and other estuarine species which inhabit, migrate through, and use areas of lower salinity for foraging, spawning, and nursery grounds. Their data imply that the impact to a given population of fish or invertebrates is highly dependent on the life stage and surrounding environmental conditions.

The significance of experimental design and exposure period on evaluating the toxicity of mercury was demonstrated in a series of studies conducted by Biesinger et al. (1982). In acute flow-through toxicity tests with *Daphnia magna*, methylmercuric chloride was about 10 times more toxic than inorganic mercury, but only about 4 times as toxic under static-renewal conditions. In the static-renewal tests with methylmercury, it was discovered that about 90 percent of the mercury had been converted to inorganic mercury during the testing period. In chronic flow-through toxicity tests with *Daphnia magna*, methylmercuric chloride was about 30 times more toxic than inorganic mercury.

Plants

Chronic toxicity (as demonstrated by reduced population growth) in a marine diatom (*Thalassiosira weissflogii*), exposed to inorganic mercury, methylmercury, dimethylmercury, and elemental mercury, was related to the aqueous concentration of a single mercury species, (the chloride species HgCl_2 or CH_3HgCl), not to the total mercury or free mercury ion concentration (Mason et al. 1996). Approximately the same concentration of CH_3HgCl and HgCl_2 reduced growth in the diatom by 50 percent. Mason et al. (1996) explain the apparently higher toxicity of methylmercury compared to inorganic mercury (expressed as a total

concentration of all inorganic forms) observed by numerous authors as a result of the low percentage of the chloride form (HgCl_2) in the inorganic mercury fraction of seawater (3.3 %) compared to the high percentage of the methyl mercuric chloride species (CH_3HgCl), which forms 100% of the methylmercury in seawater. Elemental and dimethylmercury, even though more hydrophobic than HgCl_2 and CH_3HgCl respectively, were neither accumulated nor toxic to the diatom. The hypothesis by Fisher et al. (1984), that the metals that are most bioaccumulated by phytoplankton are the most toxic, may also be true about individual mercury species.

Toxicity of Mercury in Sediment

The complex behavior of mercury in the environment makes it difficult to predict toxic effects based on bulk sediment total mercury concentrations. All the available data for effects of mercury in sediment are based on measurements of either inorganic mercury or total mercury.

The concentrations of mercury in sediment associated with toxicity are primarily derived from field studies, in contrast to the large number of laboratory toxicity tests for water exposure. The results from only two spiked sediment bioassays are available. Birge et al. (1979) reported reduced survival (70% and 45%) of rainbow trout eggs exposed to sediment contaminated with inorganic mercury (mercuric chloride) for 20 days at concentrations of 1.05 and 0.18 mg/kg respectively. Swartz et al. (1988) reported an LC50 of 13.1 mg/kg for the marine amphipod (*Rhepoxynius* sp.).

Considerable data are available, however, from field-collected samples that include both measurements of mercury concentrations in sediment and adverse biological effects. Long and MacDonald (1992) reviewed the concentrations of mercury that were associated with measures of adverse biological effects in 169 studies that included both marine and estuarine systems. Data from those studies were used to calculate Effects Range-Low (ERL) and Effects Range-Median (ERM) concentrations of 0.15 mg/kg and 0.71 mg/kg, respectively. The ERL and ERM concentrations are the lower (10 percentile) and median (50 percentile) of the study concentrations associated with toxic effects. Of the total number of studies in the data set, 8.3% had biological effects below the ERL (Long et al. 1994). The incidence of effects between the ERL and ERM concentrations was 23.5%. The incidence of effects

above the ERM concentration was 42.3% for mercury, while for other metals (e.g., chromium, copper, lead and silver) the incidence of adverse effects above the ERM was in the range of 75%. The low accuracy of the ERL and ERM mercury guidelines in predicting adverse effects compared to these other metals highlights the need for site-specific effects-based data for determining sediment mercury concentrations that are a threat to aquatic biota.

The Washington State Department of Ecology uses Apparent Effects Threshold (AET) concentrations as the basis for sediment criteria for mercury. The AETs, based on laboratory bioassays and benthic community studies, represent the concentration of a contaminant above which significant adverse effects were always observed for a specific biological indicator (PTI 1988). The AETs for mercury were empirically derived from studies conducted with contaminated and reference marine sediments collected from Puget Sound. AETs for total mercury are 0.41 mg/kg for the Microtox™ bacterial luminescence bioassay, 0.59 mg/kg for oyster larvae abnormality, 2.1 mg/kg for amphipod (*Rhepoxynius abronius*) lethality, and 2.1 mg/kg for reductions in the abundance of major taxa of benthic macroinvertebrates.

A laboratory study by McGreer (1979) demonstrated that clams, *Macoma balthica*, avoided burrowing into field-collected sediment containing a suite of metals. The concentrations of both cadmium (1.4 ppm) and mercury (0.46 ppm) best explained the behavioral responses. Avoidance of mercury-contaminated habitats by aquatic species may be important ecologically. Inhibited burrowing response, relocation, and lack of larval settlement can decrease population sizes and reduce overall community composition. Species that avoid contact with contaminated sediment and do not burrow into the sediments are more vulnerable to predators and adverse environmental conditions (e.g., temperature extremes, wave action, and contaminants in the water column).

Toxicity Associated with Mercury in Tissues

Few studies report both tissue residues and effects in either short- or long-term exposure to low concentrations of mercury (Table 2). It is important to stress that both the tissue concentration and the exposure time and route (i.e. water, food,

maternal transfer) are critical factors in producing toxic symptoms in aquatic receptors.

According to Wiener and Spry (1996), mercury transferred from the female to the eggs during oogenesis may pose a greater risk to embryos than exposure to mercury in the water column. For rainbow trout, mercury residues in ovaries of 0.5 mg/kg were associated with a significant reduction in larval survival and abnormal development (Birge et al. 1979). Whitney (1991) reported that hatching success and embryonic survival in walleye were inversely correlated with mercury concentrations in the egg (range 0.002 to 0.058 mg/kg). However, only one of 12 samples had hatching success or embryonic survival less than 90%, and there was no apparent dose-response relationship.

Mercury concentrations in brain tissue associated with lethal effects appear to show less variation than that of other tissues (e.g., muscle, whole body). For example, mercury concentrations in most types of tissues of brook trout killed by exposure to 2.9 µg/l of mercury in the water column varied among individuals, whereas concentrations in the brain showed little variation (McKim et al. 1976). These results are consistent with the hypothesis that the central nervous system, rather than muscle tissue or other organs, is the site of the most harmful toxic action in fish exposed to mercury (Wiener and Spry 1996). In their review of the literature, Wiener and Spry (1996) concluded that mercury concentrations of 7 mg/kg or greater in fish brain probably cause severe, potentially lethal effects. In sensitive species such as the walleye, brain tissue concentrations of 3 mg/kg or greater probably indicate significant toxic effects.

Based on a review of the literature, Niimi and Kisson (1994) suggest that a total mercury body burden of 1-5 mg/kg represents a threshold concentration for chronic adverse effects in aquatic organisms. Wiener and Spry (1996) reviewed the literature and provided guidance for interpreting mercury residues in the axial muscle tissue in adult fish associated with toxicity; both field and laboratory studies indicate that residues of 6 to 20 mg/kg are toxic. Whole body mercury concentrations of about 5 mg/kg in brook trout and 10 mg/kg in rainbow trout were associated with sublethal and lethal effects. Both of these papers are recent examples of attempts to identify a threshold of mercury in tissue that is associated

Table 2. Toxicity associated with mercury in tissues ($\mu\text{g/g}$ wet weight).

Fish Species	Tissue Type	Tissue Hg ($\mu\text{g/g}$, ww)	Hg Form	Exposure	Effect (s)	Reference
Oncorhynchus mykiss Rainbow trout	ovary	0.5	mercuric chloride	400-528 days	significant reduction in alevin survival (4-day post hatch); significant increase in teratogenic effects	Birge et al. 1979
Oncorhynchus mykiss Rainbow trout	whole body	10-30 30-35	methylmercuric chloride	84 days	decreased growth and appetite darkened skin and lethargy	Rodgers and Beamish 1982
Oncorhynchus mykiss Rainbow trout	brain liver muscle whole body	16-30 26-68 20-28 19	total mercury in food	270 days	darkened skin; loss in appetite, visual acuity, and growth; loss of equilibrium	Matida et al. 1971
Oncorhynchus mykiss Rainbow trout	muscle	12-23	methylmercuric chloride (4-24 ppm in food)	105 days	hyperplasia of gill epithelium	Wobeser 1975
Oncorhynchus mykiss Rainbow trout	brain liver muscle	7-32 32-114 9-52	methylmercuric chloride (4 $\mu\text{g/l}$ in water column)	30-98 days	decreased appetite and activity, mortality	Niimi and KISSOON 1994
Oncorhynchus mykiss Rainbow trout	whole body	4-27	methylmercuric chloride (9 $\mu\text{g/l}$ in water column)	12-33 days	decreased appetite and activity, mortality	Niimi and KISSOON 1994
Pimephales promelas Fathead minnow	whole body	1.4	mercuric chloride	60 days	impaired reproduction, retarded larval growth	Snarski and Olson 1982
Mugil cephalus Striped mullet	whole body	0.3	methylmercuric chloride (0.001 mg/l in water column)		inhibition of regeneration of amputated caudal fin	Weis and Weis 1978
		5.0	methylmercuric chloride (0.01 mg/l in water column)	7,11, 14 days		
Salvelinus fontinalis Brook trout	brain liver ovaries/eggs whole body	5 8 5 3	methylmercuric chloride (0.27 $\mu\text{g/l}$ in water column)	273 days	no apparent effects	McKim et al. 1976
Salvelinus fontinalis Brook trout	brain liver ovaries/eggs whole body	17 24 10 5-7	methylmercuric chloride (0.93 $\mu\text{g/l}$ in water column)	273 days	increased mortality, decreased growth, lethargy, and deformities	McKim et al. 1976
Salvelinus fontinalis Brook trout	brain liver ovaries/eggs whole body	42 58 24 24	methylmercuric chloride (2.9 $\mu\text{g/l}$ in water column)	273 days	loss of appetite, muscle spasms, and deformities; mortality	McKim et al. 1976
Stizostedion vitreum Walleye	brain liver muscle	3-6 6-14 5-8	methylmercury (5-13 ppm in food)	42-63 days	mortality; emaciation; loss of locomotion, coordination and appetite.	Scherer et al. 1975
Stizostedion vitreum Walleye	brain liver muscle	15-40 18-50 15-45	methylmercury (5-13 ppm in food)	240-314 days	higher mortality; emaciation; poorer locomotion, coordination and appetite.	Scherer et al. 1975

with adverse effects. The "thresholds" presented in these papers are based on effects in adult fish and probably do not represent a truly protective level for all species and life stages, including maternal transfer. We begin to become concerned about reproductive or early life stage effects when total Hg in whole bodies of fish are between 0.5 and 1.0 ppm.

Interactions with Other Metals

The effects on aquatic organisms due to interactions of mercury with cadmium, copper, selenium, and zinc were found to be dependent on exposure concentrations (Birge et al. 1979). In general, effects were less than additive at lower exposure concentrations and greater than additive (synergistic) at higher concentrations. Zinc and cadmium were reported to reduce the teratogenic effects of methylmercury to killifish (Weis et al. 1981). Cadmium added to methylmercury reduced the retardation effect on fin regeneration in mullet (Weis and Weis 1978).

The percentage of embryos affected and degree of malformation observed due to exposure of killifish eggs to 20-50 µg/l methylmercury was reduced when cadmium or zinc was added. Selenium was reported to reduce the developmental effects of inorganic mercury to embryos of the medaka (Japanese ricefish), but only after the formation of the embryonic liver (Bowers et al. 1980). Interactions between inorganic mercury and zinc, PCBs, and a PAH (fluoranthene) were observed to be generally additive in sediment exposure to a marine amphipod (Swartz et al. 1988). A mixture of an inorganic form of mercury (mercuric chloride) and the chlorides of zinc and lead had a synergistic toxic effect on the water exposure of a marine ciliate *Uronema marinum* (Parker 1979).

CRITERIA AND GUIDELINES

This section briefly discusses EPA's AWQC for mercury in freshwater and marine systems and various guidelines that have been proposed for evaluating the potential toxicity of mercury in sediments.

Ambient Water Quality Criteria (AWQC)

The AWQC, promulgated by the U.S. EPA, are intended neither as rules nor regulations, but present data and guidance on the effects of pollutants that can be

used to derive regulations based on considerations of water quality impacts (EPA 1993). The AWQC consist of two concentrations: the Criterion Maximum Concentration (usually referred to as the acute AWQC) and the Criterion Continuous Concentration (usually referred to as the chronic AWQC). The acute AWQC are derived from short-term toxicity tests using statistical methods that estimate the LC₅₀ concentrations for the lowest 5 percent of the most sensitive species tested. Acute AWQC for mercury are 2.4 and 2.1 µg/l for freshwater and marine organisms, respectively. They are based on inorganic mercury because it is the predominant form of mercury released into the environment (EPA 1985).

The chronic AWQC are defined by EPA as the lowest (most protective) concentrations from three categories of tests: the final chronic value, derived from chronic toxicity tests with animals; the final plant value, derived from toxicity tests using aquatic plants; and the final residue value (FRV), derived from maximum permissible tissue concentrations (for protection of human health) and bioconcentration factors (BCF). For mercury, the chronic AWQC of 0.012 µg/L for freshwater species and 0.025 µg/l for marine species represent FRVs, which are based on the Food and Drug Administration's (FDA) action level of 1 mg/kg and BCFs for methylmercury.

The use of the FDA action level to derive the chronic AWQC assumes that aquatic organisms would not be adversely affected by methylmercury tissue concentrations greater than or equal to 1 mg/kg. This is based on a study in which long-term exposure of brook trout to methylmercury resulted in tissue residue concentrations greater than 1 mg/kg but no significant effects on survival, growth, or reproduction (McKim et al. 1976). However, other studies have demonstrated that tissue concentrations close to the FDA action level may elicit adverse effects in some species (Birge et al. 1979; Snarski and Olson 1982).

Eisler (1987), in his review of the hazards of mercury to aquatic organisms, noted that the AWQC do not appear to be protective of aquatic organisms when compared to results from toxicity tests with sensitive species. Assuming the BCFs accurately reflect bioaccumulation under field conditions, the FDA action level would be the average concentration in muscle tissue of exposed aquatic organisms. Thus, many aquatic organisms exposed to mercury concentrations equivalent to

the chronic AWQC would be expected to have tissue concentrations above 1 mg/kg (EPA 1985).

Sediment Guidelines

No sediment criteria are currently available for either methylmercury or total mercury. Several approaches have been proposed for developing guidelines for screening contaminated sediments, but two of the more frequently used approaches are the National Status and Trends Effects Range approach (Long and MacDonald 1992) and the AET (PTI 1988) approach developed for screening sediments in Puget Sound, Washington. The State of Washington used the AET approach as the basis for marine sediment management standards in Puget Sound.

The ERL and ERM concentrations for mercury in marine and estuarine sediments are 0.15 and 0.71 mg/kg (Long and MacDonald 1992), respectively, while the AET concentrations for mercury range between 0.41 and 2.1 mg/kg (PTI 1988).

Tissue

No standards that would be protective of aquatic organisms have been established for mercury concentrations in tissues. The current FDA action level for the protection of human health, based only on methylmercury in the edible flesh of fish and shellfish, is 1 mg/kg (U. S. FDA 1984).

APPLICATIONS

Ecological assessments of hazardous waste sites with elevated concentrations of mercury in the aquatic environment are particularly challenging due to the complexity of the factors that affect the availability of mercury to aquatic organisms. The distribution of total mercury in sediment, which in most cases is predominantly inorganic mercury, may not provide sufficient information about the bioavailability and toxicity of mercury to aquatic species. Because of the importance of methylation in determining the availability of mercury to aquatic organisms, the sampling design, evaluation of remedial alternatives, and monitoring program should be based on an understanding of the system-specific processes that lead to increased methylation and the pathways to resources of concern. The effort required and detail of this understanding should be determined by the magnitude of the problem and the scope of the project. This section discusses some special

considerations to assist in sampling, risk assessment, monitoring, and remedial decisions.

Sampling and Monitoring Considerations

Target Species and Analysis

Mercury concentrations in resident aquatic organisms may provide the best measure of the availability of mercury in a particular area, both because of potential human health concerns and because it is the best indicator of availability of mercury under the specific conditions present at a site. In selecting target species, the trophic level, size, age, sex, life habit, metabolism, and life span of organisms are all important factors to consider. Higher trophic-level fish species are useful for determining whether a problem exists since mercury biomagnifies, and for long-term monitoring, since mercury concentrations are slower to decrease. However, even fish occupying the same trophic level, with similar diets and feeding habits, may exhibit different temporal patterns of mercury accumulation due to differences in habitat preferences, behavior, and metabolic rate causing different exposures (Jackson 1991). Mercury concentrations in biota may not correlate with sediment mercury concentrations. Correlations between mercury concentrations in predator and prey species may be useful in determining the food web pathways that connect the mercury in the sediment to the biota.

Whole body analyses of fish are typically done to determine food chain exposure, while fish fillets are typically analyzed to assess human health exposure. Whole-body mercury concentrations may be less than the concentration in the fillet; however the difference may not be statistically significant. For example, although methylmercury concentrations were higher in fillet than in whole body samples measured in four fish species (white perch, small mouth bass, bluegill, and gizzard shad) in Onondaga Lake (New York), only concentrations in bluegill were statistically different (Becker and Bigham 1995).

Invertebrates such as bivalve molluscs and mayflies (*Hexagenia* sp.) have also been used for assessing the availability of mercury in a particular location. Depending on the scope of the project, monitoring several different species from different trophic levels may be appropriate.

In developing assessment endpoints and sampling objectives, the potential for direct toxicity to aquatic organisms should not be overlooked. Laboratory and in-situ toxicity testing are useful approaches for assessing the direct biological effects of elevated mercury concentrations to aquatic organisms in sediment and water. Toxicity testing at mercury sites should include standard toxicity tests. In addition, early life-stage tests (exposure of test species from post-fertilization through embryonic, larval, and early juvenile development) or partial life-cycle tests (early juvenile through post-hatch of next generation with measurements of survival, growth, and reproductive endpoints) are sensitive tests for mercury toxicity. Monitoring changes in abundance and diversity in macrobenthic community composition may also provide useful information in assessing the toxicity of mercury in aquatic habitats.

Environmental Sampling

Investigations should be designed to include both spatial and temporal sampling. Seasonal and spatial variations in mercury concentrations, including its forms and partitioning, within a single waterbody can be significant (e.g., Gill and Bruland 1990; Parks et al. 1989). Mercury contents in mussels from different parts of the Gulf of St. Lawrence estuary (normalized by shell length and soft-tissue dry weight) were highest in areas with the greatest freshwater influence and lower in regions where the marine influence was greatest (Cossa and Rondeau 1985). Allard and Stokes (1989) found that total mercury in two species of crayfish was significantly higher in specimens from lake inlets than in those from the lake basin. Determining those environmental parameters that affect the activity of methylating microbes such as nutrients, temperature, and dissolved oxygen, and the factors affecting the availability of inorganic mercury for methylation such as the resuspension of sediment, TOC, and sulfides, may be warranted for the design of sampling and monitoring plans. Data on chloride concentration and pH may be used to determine the relative proportions of the individual inorganic and methylmercury species and their overall partition coefficients (Mason et al. 1996). In addition to the form of mercury, its partitioning between dissolved and particulate forms has an important effect on both uptake and transport. In determining the extent of contamination at a site, it is important to consider that both resuspended contaminated sediment and dissolved mercury may act as important sources.

Accurately modeling the fate of mercury in aquatic environments and the availability of mercury to aquatic organisms requires the collection of detailed information on the forms of mercury and their relative concentrations in different environmental compartments (e.g., the amounts of inorganic, methyl-, and elemental mercury in dissolved and particulate forms in the water column, sediment [particulate and pore water], and biota). The scope of a project like this is enormous and the effectiveness of this comprehensive approach has yet to be demonstrated. The transfer of mercury through the food web was modeled using a descriptive approach to explain the high levels of mercury in Lavaca Bay fish and shellfish (Evans and Engel 1994). Tissue burdens for multiple food web components are required. This approach was useful in identifying critical factors responsible for localized elevations in mercury concentrations, but also demonstrated the limitations and large effort required for modeling.

Approaches to Remediation

The level of effort needed to characterize a mercury-contaminated site may range from simple monitoring of mercury concentrations in biota and environmental media at small sites, to biogeochemical modeling at major mercury sites (PTI 1991). Although methylmercury is the form of most concern in aquatic systems, it has not been routinely measured due to the lack of standard analytical methods and cost considerations. Determining which form of mercury to measure, and in which media and organisms, depends on the nature of the contamination and the objectives of the study. Measuring only total mercury concentrations in sediment and biota may give a general picture of the extent of contamination and the magnitude of the problem, but only provides minimal information on the fate, transport, and availability of mercury in the system. In order to select an effective remedial alternative, it may be necessary to characterize the major pathways to receptors of concern and the aspects of the aquatic system that enhance methylation and influence mercury availability.

The role of speciation in determining concentrations in, and toxicity to, biota may need to be understood prior to attempts to control the geochemical cycling of mercury within a waterbody. Remediation attempts have been unsuccessful at sites where these factors have been ignored. However, this approach requires analytical

techniques that are selective and sensitive enough to measure ambient concentrations of the different mercury species.

A primary goal of many remedial investigations is to establish cleanup concentrations for mercury in various environmental media that will be protective of both human health and the environment. Establishing target cleanup concentrations for mercury is extremely difficult due to the many environmental factors that influence the transformation of inorganic mercury to methylmercury. Target cleanup concentrations should be determined on a site-by-site basis due to the variability in the bioavailability of mercury and conditions between sites. Determining a cleanup concentration requires knowing the effect threshold and translating that to a sediment concentration that is protective. Cleanup concentrations should be chosen that both reduce the source of total mercury to the system and its bioavailability to organisms. Confirmation of the effectiveness of the target cleanup concentration requires long-term monitoring of both sediment and biota.

Removing hot spots (by dredging or capping) may eliminate important sources of inorganic mercury but may not provide substantial improvement in environmental conditions if methylation rates are much higher in less contaminated areas, such as freshwater wetlands (St. Louis et al. 1994). Other factors affecting the site also need to be considered. Methylation of mercury left in sediments could be increased by dredging, increased organic loading (without sulfides), and increased thermal loading (Rada et al. 1986).

Where source control has not been feasible due to the volume and extensive distribution of the mercury, remediation strategies that focus on limiting the bioavailability of mercury have been studied. Dilution approaches, such as adding uncontaminated sediments, may reduce the supply of mercury by an order of magnitude (Rudd and Turner 1983). Complexation of mercury to “detoxify” (e.g., addition of selenium to the water column) may be a valid approach. Research indicates that selenium interferes with bioaccumulation efficiency of mercury in fish since selenium concentrates in the fish food source and excludes mercury. This approach can yield up to a twofold decrease in mercury bioaccumulation rates. However, selenium toxicity may add a new problem. Liming to increase pH in freshwater lakes has also been attempted with variable results, with positive effects

taking two or more years (Lindqvist 1991; Meili 1995). The long-term effectiveness of these remedies and the potential adverse effects on the environment have not been well-studied.

Analytical Considerations

Detection Limits in Water, Sediment, and Biota

Detection limits should be chosen based on the objective of the study. Analysis of mercury in water samples is particularly difficult due to the very low concentrations (parts per trillion) that need to be measured. Achieving low detection limits is further complicated by the possibility of external contamination of samples which can be a significant problem (Fitzgerald 1990).

The chronic AWQC for methylmercury are 0.012 and 0.025 $\mu\text{g/l}$ for freshwater and saltwater, respectively. Using the AWQC as a detection limit may be difficult as few labs have analytical procedures that can reach these low concentrations. The detection limit for the EPA standard contract lab program (CLP) method for analysis of total mercury in water is 0.2 $\mu\text{g/l}$. To achieve ecologically relevant detection limits it may be necessary to employ analytical methodologies other than those specified under the CLP, or to modify the CLP methods. For example, mercury detection limits to determine the mass balance of mercury in the Onondaga Lake Superfund site were established as follows: 0.00001 $\mu\text{g/l}$ for methylmercury in water; 0.0001 $\mu\text{g/l}$ for total mercury in water; and 0.01 mg/kg wet weight in fish (PTI 1991).

These detection limits were achievable by modifying standard procedures. Unfiltered water samples are analyzed for comparison to criteria concentrations. In the EPA criteria document, the measurement of acid-soluble mercury (the mercury that passes through a 45- μm filter following acidification to a pH of 1.5 to 2.0) is recommended, although no EPA-approved protocol has been established (EPA 1985). Some recent analytical approaches for the analysis of methylmercury are described by the following authors: Bloom and Crecelius (1983), Gill and Fitzgerald (1987), Bloom (1989), Gill and Bruland (1990).

Choosing appropriate detection limits for sampling of sediment is particularly important for mercury sites because even low concentrations can cause significant

accumulations in biota. Detection limits for sediments should be below the ERL value of 0.15 mg/kg developed by Long and MacDonald (1992).

Quality control is an important aspect of any testing program but is particularly important for the analysis of mercury in environmental samples. It is highly recommended that analyses of mercury in water, sediment, and tissue also include analyses of certified standards for the appropriate matrix as part of the quality control plan to verify the extraction and analytical processes.

Difficulties in the extraction and analysis of mercury residues in tissues are apparently not uncommon. For example, in three recent Superfund projects in three different regions, methylmercury concentrations in fish tissue were reported to be higher than total mercury concentrations. In another study of contaminant concentrations in tissues of aquatic organisms, spike-recovery values for mercury in tissue samples were in the range of 50 percent (Tetra Tech 1988).

The National Research Council of Canada has certified standards for methylmercury and total mercury in animal tissue samples (e.g., dogfish liver and dogfish muscle), and for sediment and water samples. The U.S. National Bureau of Standards has comparable standards for total mercury.

SUMMARY

NOAA recommends a site-specific approach that focuses on determining the availability of mercury and the potential for toxic effects. The accumulation of mercury in aquatic biota is often the primary concern at mercury sites and is useful for assessing availability. Bioaccumulation studies should measure tissue concentrations in more than one resident and/or transplanted caged species, preferably with species representing different trophic levels or different food web pathways. It may not be possible to correlate sediment mercury concentrations with concentrations in biota. However, correlations between mercury concentrations in predator and prey species may be useful in determining pathways of mercury transfer.

Toxicity tests such as the standard amphipod tests should also be conducted to assess mercury toxicity to aquatic organisms. At major mercury sites, chronic

toxicity endpoints should be included in the assessment—in particular, fish early life stage or reproductive endpoint tests. Because of the persistence of mercury in aquatic systems, source control alone may not be sufficient to permit recovery. Additional remedial actions may be required to reduce the total mercury burden in the system. Long-term monitoring of tissue concentrations of mercury in aquatic biota is needed to assess remedial effectiveness.

REFERENCES

- Alberts, J.J., M.T. Price and M. Kania. 1990. Metal concentrations in tissues of *Spartina alterniflora* (Loisel.) and sediments of Georgia salt marshes. *Estuarine, Coastal and Shelf Science* 30: 47-58.
- Allard, M. and P. Stokes. 1989. Mercury in crayfish species from thirteen Ontario Lakes in relation to water chemistry and smallmouth bass (*Micropterus dolomieu*) mercury. *Canadian Journal of Fisheries and Aquatic Sciences* 46.
- Andersen, A., K. Julshamn, O. Ringdal, and J. Morkore. 1987. Trace Elements Intake in the Faroe Islands, II: Intake of Mercury and Other Elements by Consumption of Pilot Whales (*Globicephalus meleanus*). *Science of the Total Environment* 65:63-68.
- Andre, J. M., A. Boudou, and F. Ribeyre, and M. Bernhard. 1991a. Comparative study of mercury accumulation in dolphins (*Stenella coeruleoalba*) from French Atlantic and Mediterranean coasts. *Sci. Total Environ.* 104:191-209.
- Andre, J. M., A. Boudou, and F. Ribeyre. 1991b. Mercury accumulation in Delphinidae. *Wat. Air Soil Pollut.* 56:187-201.
- Arima, S., and K. Nagakura. 1979. Mercury and selenium content of *Odontoceti*. *Bull. Japanese Soc. Scientific Fish.* 45(5):623-626.
- Armstrong, F. A. J. 1979. Effects of mercury compounds on fish. In: J. O. Nriagu (Ed.), *The Biogeochemistry of Mercury in the Environment*. pp. 657-670. New York: Elsevier/North-Holland Biomedical Press.

- Battelle. 1987. Measurement of bioavailable mercury species in fresh water and sediments. Palo Alto, California: Electric Power Research Institute.
- Becker, D.S. and G.N. Bigham. 1995. Distribution of mercury in the aquatic food web of Onondaga Lake, New York. *Water, Air, and Soil Pollution* 80:563-571.
- Becker, D. S., G. N. Bigham, and M. H. Murphy. 1993. Distribution of mercury in a lake food web. Poster presented at 14th annual meeting of the Society of Environmental Toxicology and Chemistry (SETAC), Houston, Texas, November 14-18, 1993.
- Beijer, K. and A. Jernelov. 1979. Methylation of mercury in aquatic environments. In: J. O. Nriagu (Ed.), *The Biogeochemistry of Mercury in the Environment*. pp. 203-210. New York: Elsevier/North-Holland Biomedical Press.
- Benes, P. and B. Havlik. 1979. Speciation of mercury in natural waters. In: J. O. Nriagu (Ed.), *The Biogeochemistry of Mercury in the Environment*. pp. 175-202. New York: Elsevier/North-Holland Biomedical Press.
- Berman, M. and R. Bartha. 1986. Levels of chemical versus biological methylation of mercury in sediments. *Bulletin of Environmental Contamination and Toxicology* 36:401-404.
- Biesinger, K.E., L.E. Anderson, and J.G. Eaton. 1982. Chronic effects of inorganic and organic mercury on *Daphnia magna*: toxicity, accumulation, and loss. *Arch. Environ. Contam. Toxicol.* 11: 769-774.
- Birge, W. J., J. A. Black, A. G. Westerman, and J. E. Hudson. 1979. The effects of mercury on reproduction of fish and amphibians. In: J. O. Nriagu (ed.). *The Biogeochemistry of Mercury in the Environment*. pp. 629-655. New York: Elsevier/North-Holland Biomedical Press.
- Bjerregaard, P. and L. Christensen. 1993. Accumulation of organic and inorganic mercury from food in the tissues of *Carcinus maenas*: effect of waterborne selenium. *Marine Ecol. Prog. Ser.* 99: 271-281.
- Bjornberg, A., L. Håkanson, and L. Lundbergh. 1988. A theory on the mechanisms regulating the bioavailability of mercury in natural waters. *Environmental Pollution* 49:53-61.

Bloom, N.S. 1989. Determination of picogram levels of methylmercury by aqueous phase ethylation, followed by cryogenic gas chromatography with cold vapor atomic fluorescence detection. *Canadian Journal of Fisheries and Aquatic Sciences* 46:1131-1140.

Bloom, N. S. 1992. On the chemical form of mercury in edible fish and marine invertebrate tissue. *Can. J. Fish. Aquat. Sci.* 49:1010-1017.

Bloom, N.S., C.J. Watras, and J.P. Hurley. 1991. Impact of acidification on the methylmercury cycle of remote seepage lakes. *Water, Air, Soil Pollut.* 56:477-491.

Bloom, N.S. and S.W. Effler. 1990. Seasonal variability in the mercury speciation of Onondaga Lake (New York). *Water, Air, Soil Pollut.* 53:251-265.

Bloom, N.S. and E.A. Crecelius. 1983. Determination of mercury in seawater at subnanogram per litre levels. *Marine Chemistry* 14:49-59.

Blum, J. M. and R. Bartha. 1980. Effect of salinity on methylation of mercury. *Bulletin of Environmental Contamination and Toxicology* 25:404-408.

Bonzongo, J.J., Heim, K.J., Y. Chen, W.B. Lyons, J.J. Warwick, G.C. Miller and P.J. Lechler. 1996. Mercury pathways in the Carson River-Lahontan Reservoir System, Nevada, USA. *Environ. Toxicol. and Chem.* 15(5):677-683.

Born, E. W., I. Kraul, and T. Kristensen. 1981. Mercury, DDT, and PCB in the Atlantic walrus (*Odobenus rosmarus rosmarus*) from the Thule District, North Greenland. *Arctic* 34:255.

Boudou, A., M. Delnomdedieu, D. Georgescauld, F. Ribeyre, and E. Saouter. 1991. Fundamental roles of biological barriers in mercury accumulation and transfer in freshwater ecosystems (analysis at organism, organ, cell and molecular levels). *Water, Air, and Soil Pollution* 56:807-822.

Bowers, M. A., D. Dostal, and J. F. Heisinger. 1980. Failure of selenite to protect against mercuric chloride in early developmental stages of the Japanese ricefish (*Oryzias latipes*). *Comparative Biochemistry and Physiology* 66C:175-178.

- Breteler, R.J., I. Valiela, and J.M. Teal. 1981. Bioavailability of mercury in several North-eastern U.S. *Spartina* ecosystems. *Estuarine, Coastal and Shelf Science* 12: 155-166.
- Brock, V. 1993. Effects of mercury on the physiological condition and content of the biomarker ALA in the oyster *Ostrea edulis*. *Mar. Ecol. Prog. Series*. 96:169-175.
- Brown, D. A., R. W. Gosset, P. Hershelman, H. A. Schaefer, K. D. Jenkins, and E. M. Perkins. 1983. Bioaccumulation and detoxification of contaminants in marine organisms from Southern California coastal waters. In: D. F. Soule and D. Walsh (Eds.), *Waste Disposal in the Oceans*. p. 171. Boulder: Westview Press.
- Bryan, G. W. and W. J. Langston. 1992. Bioavailability, accumulation and effects of heavy metals in sediments with special reference to United Kingdom estuaries: a review. *Environ. Pollut.* 76: 89-131.
- Calabrese, A., J. R. MacInnes, D. A. Nelson, and J. E. Miller. 1977. Survival and growth of bivalve larvae under heavy metal stress. *Mar. Biol.* 41:179-184.
- Callister, S. M. and Winfrey, M. R. 1986. Microbial methylation of mercury in upper Wisconsin River sediments. *Water, Air and Soil Pollution* 29: 453-465.
- Choi, S.-C., and R. Bartha. 1994. Environmental factors affecting mercury methylation in estuarine sediments. *Bull. Environ. Contam. Toxicol.* 53:805-812.
- Clarkson, T. W. 1972. Recent advances in toxicology of mercury with emphasis on the alkyl mercurials. *Crit. Rev. Toxicol.* 203-234:.
- Compeau, G. and R. Bartha. 1983. Effects of sea salt anions on the formation and stability of methylmercury. *Bulletin of Environmental Contamination and Toxicology* 31:486-493.
- Compeau, G. and R. Bartha. 1985. Sulfate-reducing bacteria: principal methylators of mercury in anoxic estuarine sediment. *Appl. Environ. Microbiol.* 50: 498-502.
- Cope, W. G., J. G. Wiener, and R. G. Rada. 1990. Mercury accumulation in yellow perch in Wisconsin seepage lakes: relation to lake characteristics. *Environmental Toxicology and Chemistry* 9:931-940.

- Cossa, D. and J. G. Rondeau. 1985. Seasonal, geographical and size-induced variability in mercury content of *Mytilus edulis* in an estuarine environment: a re-assessment of mercury pollution level in the Estuary and Gulf of St. Lawrence. *Marine Biology* 88:43-49.
- Craig, P.J., and P.A. Moreton. 1983. Total mercury, methyl mercury and sulphide in River Carron sediments. *Marine Pollution Bulletin* 14(11):408-411.
- Craig, P.J., and P.A. Moreton. 1985. The role of speciation in mercury methylation in sediments and water. *Environmental Pollution Series B* 10:141-158.
- Darnell, D. W., B. Greene, M. T. Henzl, J. M. Hosea, R. A. McPherson, J. Sneddon, M. D. Alexander. 1986. Selective recovery of gold and other metal ions from an algal biomass. *Environ. Sci. Technol.* 20: 206-208.
- De Haan, H. 1992. Impacts of environmental changes on the biogeochemistry of aquatic humic substances. *Hydrobiologia* 229:59-71.
- Dukerschein, J. T., R. G. Rada, and M. T. Steingraeber. 1992. Cadmium and mercury in emergent mayflies (*Hexagenia bilineata*) from the upper Mississippi River. *Arch. Environ. Contam. Toxicol.* 23: 109-116.
- Eganhouse, R.P. and D.R. Young. 1978. Total and organic mercury in benthic organisms near a major submarine wastewater outfall system. *Bulletin of Environmental Contamination and Toxicology* 19:758-766.
- Eisler, R. 1987. Mercury hazards to fish, wildlife, and invertebrates: a synoptic review. U.S. Fish and Wildlife Service Biology Report 85(1.10). Washington, D.C.: U.S. Department of the Interior. 90pp.
- Evans, D.W. and D.W. Engel. 1994. Mercury bioaccumulation in finfish and shellfish from Lavaca Bay, Texas: descriptive models and annotated bibliography. NOAA Technical Memorandum NMFS-SEFSC-348. Beaufort, North Carolina: National Marine Fisheries Service, NOAA.
- Faust, B.C. 1992. The octanol/water distribution coefficients of methylmercuric species: the role of aqueous-phase chemical speciation. *Environ. Toxicol. Chem.* 11:1373-1376.

- Feltier, J. S., E. Kahn, B. Salick, F. C. Van Natta, and M. W. Whitehouse. 1972. *Ann. Intern. Med.* 76:779-792.
- Fisher, N.S., M. Bohé, and J.-L. Teyssié. 1984. Accumulation and toxicity of Cd, Zn, Ag, and Hg in four marine phytoplankters. *Mar. Ecol. Prog. Ser.* 18:201-213.
- Fitzgerald, W.F. 1990. Mercury in seawater. In: *Mercury in the Marine Environment. Workshop proceedings.* U.S. Dept. Interior, Mineral Management Service, Anchorage, Alaska. OCS Study MMS 89-0049.
- Fowler, S. W., M. Heyraud, and J. La Rosa, 1978. Factors affecting methyl and inorganic mercury dynamics in mussels and shrimp. *Marine Biology* 46: 267-276.
- Francesconi, K. and R. C. J. Lenanton 1992. Mercury contamination in a semi-enclosed marine embayment: organic and inorganic mercury content of biota, and factors influencing mercury levels in fish. *Marine Environmental Research* 33: 189-212.
- Gadd, G. M. 1988. Accumulation of metals by microorganisms and algae. In: Rehm H-J and G. Reed (eds) *Biotechnology*. Vol 6b, 401-433. Weinheim, Germany: VCH Verlagsgesellschaft.
- Gentile, J. H., S. M. Gentile, G. Hoffman, J. F. Heltshe, and N. Hairston, Jr. 1983. The effects of a chronic mercury exposure on survival, reproduction and population dynamics of *Mysidopsis bahia*. *Environ. Toxicol. Chem.* 2:61-68.
- Giblin, F. J. and E. J. Massaro. 1973. Pharmacodynamics of methyl mercury in rainbow trout (*Salmo gairdneri*): tissue uptake, distribution and excretion. *Toxicol. Appl. Pharmacol.* 24:81-91.
- Gill, G. A. and K. W. Bruland. 1990. Mercury speciation in surface freshwater systems in California and other areas. *Environmental Science and Technology* 24(9): 1392-1400.
- Gill, G.A. and W.F. Fitzgerald. 1987. Picomolar mercury measurements in seawater and other materials using stannous chloride reduction and two-stage gold amalgamation with gas phase detection. *Marine Chemistry* 20: 227-243.

- Gilmour, C.C., E.A. Henry, and R. Mitchell. 1992. Sulfate stimulation of mercury methylation in freshwater sediments. *Environ. Sci. Technol.* 26(11): 2281-2287.
- Gilmour, C. C. and E. A. Henry. 1991. Mercury methylation in aquatic systems affected by acid deposition. *Environmental Pollution* 71(2-4): 131-169.
- Gilmour, C. C. and Capone, D. G. 1987. Relationship between Hg methylation and the sulfur cycle in estuarine sediments. *EOS* 68:1718.
- Greib, T.M., C.T. Driscoll, S. P. Gloss, C.L. Schofield, G.L. Bowie, and D.B. Porcella. 1990. Factors affecting mercury accumulation in fish in the upper Michigan peninsula. *Environmental Toxicology and Chemistry* 9:919-930.
- Hall, A.S., F.M. Teeny, and E.J. Gauglitz. 1976a. Mercury in fish and shellfish of the northeast Pacific. II. Sablefish, *Anoplopoma fimbria*. *Fishery Bulletin* 74:791-797.
- Hall, A.S., F.M. Teeny, L.G. Lewis, W.H. Hardman, and E.J. Gauglitz, Jr. 1976b. Mercury in fish and shellfish of the northeast Pacific. I. Pacific Halibut, *Hippoglossus stenolepis*. *Fishery Bulletin* 74:783-789.
- Hansen, C. T., C. O. Nielsen, R. Dietz, and M. M. Hansen. 1990. Zinc, cadmium, mercury and selenium in Minke Whales, Belugas and Narwhals from West Greenland. *Polar Biology* 10:529-539.
- Hansen, D. J. 1989. U.S. Environmental Protection Agency regulations and criteria for mercury in water. Summary presentation to the coordination team. In: *Mercury in the Marine Environment. Workshop proceedings.* U.S. Dept. Interior, Mineral Management Service, Anchorage, Alaska. OCS Study MMS 89-0049.
- Hara, T.J., Y.M.C. Law, and S. McDonald. 1976. Effects of mercury and copper on the olfactory response in rainbow trout (*Oncorhynchus mykiss*). *J Fish Res Bd Can* 33:1568-1573.
- Heisinger, F. J. and W. Green. 1975. Mercuric chloride uptake by eggs of the rice fish and resulting teratogenic effects. *Bull. Environ. Contam. Toxicol.* 14:665-673.
- Hill, W.R., A.J. Stewart, and G.E. Napolitano. 1996. Mercury speciation and bioaccumulation in lotic primary producers and primary consumers. *Can. J. Fish. Aquat. Sci.* 53:812-819.

- Hines, M. E., S. L. Knollmeyer, and J.B. Tugel. 1989. Sulfate reduction and other sedimentary biogeochemistry in a northern New England salt marsh. *Limnol. Oceanogr.* 34(3): 578-590.
- Hintelmann, H., P. M. Welbourn, and R. D. Evans. 1995. Binding of methylmercury compounds by humic and fulvic acids. *Water, Air, and Soil Pollution* 80:1031-1034.
- Hoff, R., H. Curl, Jr., J. Farr, and N. Beckvar. 1994. *Empire Knight: Assessing environmental risk*. NOAA Technical Memorandum NOAA ORCA 81. Seattle: Hazardous Materials Response and Assessment Division, National Oceanic and Atmospheric Administration. 31pp.
- Holden, A. V. 1978. Pollutants and seals-A review. *Mammal Rev.* 8(1-2):53-66.
- Horvat, M. 1991. Determination of methylmercury in biological certified reference materials. *Water, Air, and Soil Pollution* 56:95-102.
- Huckabee, J., J. Elwood, and S. Hildebrand. 1979. Accumulation of mercury in freshwater biota. In: Nriagu (ed.) *The Biogeochemistry of Mercury in the Environment*. pp 277-302. New York: Elsevier/North-Holland Biomedical Press 1979.
- Hughes, W. L. 1957. A physicochemical rationale for the biological activity of mercury and its compounds. *Ann. New York Acad. Sci.* 65: 454-460.
- Jackson, T.A. 1991. Biological and environmental control of mercury accumulation by fish in lakes and reservoirs of northern Manitoba, Canada. *Can. J. Fish. Aquat. Sci.* 48:2449-2470.
- Jackson, T. A. 1986. Methyl mercury levels in a polluted prairie river-lake system: seasonal and site specific variations, and the dominant influence of trophic conditions. *Canadian Journal of Fisheries and Aquatic Sciences* 43:1873-1877.
- Jackson, T. A. 1987. Methylation, demethylation, and bio-accumulation of mercury in lakes and reservoirs of northern Manitoba, with particular reference to effects of environmental changes caused by the Churchill-Nelson River diversion. In: *Summary Report, Canada-Manitoba Agreement on the Study and Monitoring of Mercury in the Churchill River Diversion*. Ottawa: Governments of Canada and Manitoba.

- Jackson, T. A. 1988. Accumulation of mercury by plankton and benthic invertebrates in riverine lakes of northern Manitoba (Canada): Importance of regionally and seasonally varying environmental factors. *Canadian Journal of Fisheries and Aquatic Sciences* 45:1744-1757.
- Jackson, T. A. 1989. The influence of clay minerals, oxides, and humic matter on the methylation and demethylation of mercury by micro-organisms in freshwater environments. *Applied Organometallic Chemistry* 3:1-30.
- Jernelov, A. 1968. Laboratory Experiments regarding the conversion of mercury into its different forms of occurrence 1. *Vatten* 24(1):53-56. Translation by Canada Dept. Sed. State Transl. Bur..
- Joiris, C. R., L. Holsbeek, J. M. Bouquegneau, and M. Bossicart. 1991. Mercury contamination of the harbour porpoise *Phocoena phocoena* and other cetaceans from the North Sea and the Kattegat. *Wat. Air Soil Pollut.* 56: 283-293.
- Julshamn, K. O. Ringdal, and O. R. Braekkan. 1982. Mercury concentration in liver and muscle of Cod (*Gadus morhua*) as an evidence of migration between waters with different levels of mercury. *Bull. Environm. Contam. Toxicol.* 29:544-549.
- Julshamn, K., A. Andersen, O. Ringdal, and J. Morkore. 1987. Trace elements intake in the Faroe Islands, I. Element levels in edible parts of Pilot Whales (*Globicephalus meleanus*). *Sci. Total Environ.* 65:53-62.
- Kelly, C.A., J.W.M. Rudd, V.L. St. Louis, and A. Heyes. 1995. Is total mercury concentration a good predictor of methyl mercury concentration in aquatic systems? *Water, Air, and Soil Pollution* 80:715-724.
- Kihlstrom, J. E. and L. Hulth. 1972. The effect of phenylmercuric acetate upon the frequency of hatching in the zebrafish. *Bull. Environ. Contam. Toxicol.* 7:111.
- Kobayashi, N. 1984. Marine ecotoxicological testing with echinoderms. In: G. Persoone, E. Jaspers, and C. Claus (Eds.), *Ecotoxicological Testing for the marine Environment*. Bredene, Belgium: State University Ghent and Institute Mar. Scient. Res. pp. 341-381.

- Kopfler, F. C. 1974. The accumulation of organic and inorganic mercury compounds by the eastern oyster (*Crassostrea virginica*). *Bull. Environ. Contam. Toxicol.* 11:275-280.
- Korthals, E. T and M. R. Winfrey. 1987. Seasonal and spatial variations in mercury methylation and demethylation in an oligotrophic lake. *Applied and Environmental Microbiology* 53:2397-2404.
- Langston, W. J. 1990. Chapter 7: Toxic effects of metals and the incidence of metal pollution in marine environments. In: R. W. Furness and P. S. Rainbow (Eds.), *Heavy Metals in the Marine Environment*. pp. 101-122. Boca Raton, Florida: CRC Press, Inc.
- Langston, W. J. 1986. Metals in sediments and benthic organisms in the Mersey Estuary. *Estuarine, Coastal and Shelf Science* 23:239-261.
- Langston, W. J. 1982. The distribution of mercury in British estuarine sediments and its availability to deposit-feeding bivalves. *J. Mar. Biol. Ass. U.K.* 62: 667-684.
- Lasorsa, B. and S. Allen-Gil. 1995. The methylmercury to total mercury ratio in selected marine, freshwater, and terrestrial organisms. *Water, Air, and Soil Pollution* 80:905-913.
- Leah, R.T., S.J. Evans, and M.S. Johnson. 1992. Mercury in flounder (*Platichthys flesus* L.) from estuaries and coastal waters of the north-east Irish Sea. *Environmental Pollution* 75:317-322.
- Lenka, M., K.K. Panda, and B. B. Panda. 1990. Studies on the ability of water hyacinth (*Eichlornia crassipes*) to bioconcentrate and biomonitor aquatic mercury. *Environ. Pollut.* 66: 89-99.
- Leonzio, C., S. Focardi, and C. Fossi. 1992. Heavy metals and selenium in stranded dolphins of the Northern Tyrrhenian (NW Mediterranean). *Sci. Total Environ.* 119:77-84.
- Lindberg, S. E. and R.C. Harriss. 1974. Mercury-organic matter associations in estuarine sediments and interstitial water. *Envir. Sci. Tech.* 8(5):459-462.

Lindqvist, O. ed. 1991. Mercury in the Swedish environment: Recent research on causes, consequences and corrective methods. *Water, Air, and Soil Pollution* 55(1-2). Special Issue. 261pp.

Long, E. R., D. D. MacDonald, S. L. Smith, and F. D. Calder. 1995. Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments. *Environmental Management* 19(1):81-97.

Long, E. R. and D. D. MacDonald. 1992. National Status and Trends Program Approach. In: *Sediment Classification Methods Compendium*. EPA 823-R-92-006. EPA Office of Water (WH-556). Washington, DC.: U.S. Environmental Protection Agency.

Luoma, S.N. 1977. The dynamics of biologically available mercury in a small estuary. *Estuarine Coast. Mar. Sci.* 5:643-652.

Major, M.A., D.H. Rosenblatt and K.A. Bostian. 1991. The octanol/water partition coefficient of methylmercuric chloride and methylmercuric hydroxide in pure water and salt solutions. *Environ. Toxicol. Chem.* 10:5-8.

Mason, R.P., J.R. Reinfelder, and F.M.M. Morel. 1996. Uptake, toxicity, and trophic transfer of mercury in a coastal diatom. *Environ. Sci. Technol.* 30:1835-1845.

Mason, R.P., K.R. Rolfhus and W.F. Fitzgerald. 1995a. Methylated and elemental mercury cycling in surface and deep ocean waters of the North Atlantic. *Water, Air, and Soil Pollution* 80: 665-677.

Mason, R.P., J.R. Reinfelder, and F.M.M. Morel. 1995b. Bioaccumulation of mercury and methylmercury. *Water, Air, and Soil Pollution* 80:915-921.

Mathers, R. and P. Johansen. 1985. The effects of feeding ecology on mercury accumulation in walleye (*Stizostedion vitreum*) and pike (*Esox lucius*) in Lake Simcoe. *Canadian Journal of Zoology* 63: 2006-2012.

Matida, Y., H. Kumada, S. Kimura, Y. Saiga, T. Nose, M. Yokote, and H. Kawatsu. 1971. Toxicity of mercury compounds to aquatic organisms and accumulation of the compounds by the organisms. *Bull. Freshwater Fish. Res. Lab. (Tokyo)* 21:197-227.

- McGreer, E. R. 1979. Sublethal effects of heavy metal contaminated sediments on the bivalve *Macoma balthica* (L). *Mar. Pollut. Bull.* 10:259.
- McKenney, C. L., Jr. and J. D. Costlow, Jr. 1981. The effects of salinity and mercury on developing megalopae and early crab stages of the blue crab *Callinectes sapidus* Rathbun. In: F. J. Vernberg, A. Calabrese, F. P. Thurberg, and W. B. Vernberg (Eds.), *Biological Monitoring of Marine Pollutants*. pp. 241-262. New York: Academic Press.
- McKim, J.M., G. F. Olson, G. W. Holcombe, and E.P. Hunt. 1976. Long-term effects of methylmercuric chloride on three generations of brook trout (*Salvelinus fontinalis*): Toxicity, accumulation, distribution, and elimination. *Journal Fisheries Research Board of Canada* 33:2726-2739.
- Meili, M.. 1995. Liming effects on mercury concentrations in fish. In: L. Henrikson and Y. W. Brodin (eds). *Liming Acidified Sur. Waters*. p. 383-398. Berlin: Springer.
- Mierle, G., and R. Ingram. 1991. The role of humic substances in the mobilization of mercury from watersheds. *Water, Air, and Soil Pollution* 56:349-357.
- Miskimmin, B. M., J. W. M. Rudd, and C. Kelly. 1992. Influence of dissolved organic carbon, pH, and microbial respiration rates on mercury methylation and demethylation in lake water. *Can. J. Fish. Aquat. Sci.* 49:17-22.
- Møhlenberg, F. and H.U. Riisgård. 1988. Partitioning of inorganic and organic mercury in cockles *Cardium edule* (L.) and *C. Glaucom* (Bruguere) from a chronically polluted area: influence of size and age. *Environ. Pollut.* 55:137-148.
- Nicoletto, P. and A. Hendricks. 1987. Sexual differences in accumulation of mercury in four species of centrarchid fishes. *Canadian Journal of Zoology* 66:944-949.
- Niimi, A.J. and G.P. Kisson. 1994. Evaluation of the critical body burden concept based on inorganic and organic mercury toxicity to rainbow trout (*Oncorhynchus mykiss*). *Arch. Environ. Contam. Toxicol.* 26:169-178.
- Nishimura H., and M. Kumagai. 1983. Mercury pollution of fishes in Minamata Bay and surrounding water: analysis of pathway of mercury. *Water, Air, and Soil Pollution* 20:401-411.

Olson, B.H. and R.C. Cooper. 1976. Comparison of aerobic and anaerobic methylation of mercuric chloride by San Francisco Bay sediments. *Water Res.* 10: 113-116.

Olson, K R., K. S. Squibb, and R. J. Cousins. 1978. Tissue uptake, subcellular distribution, and metabolism of $^{14}\text{CH}_3\text{HgCl}$ and $\text{CH}_3^{203}\text{HgCl}$ by rainbow trout, *Salmo gairdneri*. *J. Fish. Res. Board Can.* 35:381-390.

Parks, J. W., A. Lutz, and J. A. Sutton. 1989. Water column methylmercury in the Wabigoon/English River-Lake System: Factors controlling concentration, speciation, and net production. *Canadian Journal of Fisheries and Aquatic Sciences* 46:2184-2202.

Passow, H., A. Rothstein, and T. Clarkson. 1961. The general pharmacology of the heavy metals. *Pharmacol. Rev.* 13:185-224.

Pentreath, R.J. 1976a. The accumulation of organic mercury from sea water by the plaice, *Pleuronectes platessa* L. *Journal of Experimental Marine Biology and Ecology* 24:121-132.

Pentreath, R.J. 1976b. The accumulation of inorganic mercury from sea water by the plaice, *Pleuronectes platessa* L. *Journal of Experimental Marine Biology and Ecology* 24:103-119.

PTI. 1991. Onondaga Lake RI/FS Work Plan. PTI Environmental Services, Bellevue, Washington.

PTI. 1988. Briefing report to the EPA Science Advisory Board: The Apparent Effects Threshold approach. Seattle: Environmental Protection Agency, Region 10, Office of Puget Sound. 57 pp.

Rada, R. G., J. G. Wiener, M. R. Winfrey, D. E. Powell. 1989a. Recent increases in atmospheric deposition of mercury to north-central Wisconsin lakes inferred from sediment analyses. *Arch. Environ. Contam. Toxicol.* 18:175-181.

Rada, R. G., J. E. Findley, and J. G. Wiener. 1989b. Environmental fate of mercury discharged into the upper Wisconsin River. *Water, Air, and Soil Pollution* 29:57-76.

- Rada, R. G., J. E. Findley, and J. G. Wiener. 1986. Environmental fate of mercury discharged into the upper Wisconsin River. *Water Air Soil Pollut.* 29:57-76.
- Ramlal, P. S., J.W.M. Rudd, A. Furutani, L. Xun. 1985. The effect of pH on methyl mercury production and decomposition in lake sediments. *Canadian Journal of Fisheries and Aquatic Sciences* 42: 685-692.
- Regnell, O. 1994. The effect of pH and dissolved oxygen levels on methylation and partitioning of mercury in freshwater model systems. *Environmental Pollution* 84:7-13.
- Rehnberg, B.C. and C.B. Schreck. 1986. Acute metal toxicology of olfaction in coho salmon: behavior, receptors, and odor-metal complexation. *Bull. Environ. Contam. Toxicol.* 36:579-586.
- Reigel, D. V. 1990. The distribution and behavior of mercury in sediments and marine organisms of Lavaca Bay, Texas. College Station, Texas: The Texas A&M University. Masters Thesis.
- Richardson, G.M. and D.J. Currie. 1996. Does acid precipitation exacerbate the problem of fish mercury contamination? *SETAC News* 16(2): 14.
- Riisgård , H.U. and P. Famme. 1988. Distribution and mobility of organic and inorganic mercury in flounder, *Platichthys flesus*, from a chronically polluted area. *Toxicol. Environ. Chem.* 16:219-228.
- Riisgård , H. U. and P. Famme. 1986. Accumulation of inorganic and organic mercury in shrimp, *Crangon crangon*. *Marine Pollution Bulletin* 17:255-257.
- Riisgård, H.U., T. Kiorboe, F. Møhlenberg, I. Drabk, and P. Pheiffer Madsen. 1985. Accumulation, elimination and chemical speciation of mercury in the bivalves *Mytilus edulis* and *Macoma balthica*. *Mar. Biol.* 86:55-62.
- Riisgård , H. U. and S. Hansen. 1990. Biomagnification of mercury in a marine grazing food-chain: algal cells *Phaeodactylum tricorutum*, mussels *Mytilus edulis* and flounders *Platichthys flesus* studied by means of a stepwise-reduction-CVAA method. *Marine Ecology Progress Series* 62:259-270.

- Rodgers, D.W. and F.W.H. Beamish. 1982. Dynamics of dietary methylmercury in rainbow trout, *Salmo gairdneri*. *Aquatic Toxicology* 2: 271-290.
- Rodgers, D.W. and F. W. H. Beamish. 1981. Uptake of waterborne methylmercury by rainbow trout (*Salmo gairdneri*) in relation to oxygen consumption and methylmercury concentration. *Canadian Journal of Fisheries and Aquatic Sciences* 38(11): 1309-1315.
- Roesijadi, G., A.S. Drum, and J.R. Bridge. 1981. Mercury in mussels of Bellingham Bay, Washington (U.S.A.): the occurrence of mercury-binding proteins. In: Vernberg, F.J., Calabrese, A. Thurberg, F.P. and W.B. Vernberg (Eds.), *Biological Monitoring of Marine Pollutants*. pp. 357-376. New York: Academy Press.
- Rubinstein, N.I., E. Loes, and N.R. Gregory. 1983. Accumulation of PCBs, mercury and cadmium by *Nereis virens*, *Mercenaria mercenaria* and *Palaemonetes pugio* from contaminated harbor sediments. *Aquatic Toxicology* 3:249-260.
- Rudd, J. W. M. and M. A. Turner. 1983. The English-Wabigoon River System: V. Mercury and selenium bioaccumulation as a function of aquatic primary productivity. *Canadian Journal of Fisheries and Aquatic Sciences* 40:2251-2259.
- Rudd, J. W. M., M. A. Turner, A. Furutani, A. L. Swick, and B. E. Townsend. 1983. The English-Wabigoon River System: I. A synthesis of recent research with a view towards mercury amelioration. *Canadian Journal of Fisheries and Aquatic Sciences* 40:2206-2217.
- Saouter, E., L. Hare, P. G. C. Campbell, A. Boudou, and F. Ribeyre. 1993. Mercury accumulation in the burrowing mayfly *Hexagenia rigida* (Ephemeroptera) exposed to CH_3HgCl or HgCl_2 in water and sediment. *Wat. Res.* 27(6):1041-1048.
- Saouter, E. F. Ribeyre, A. Boudou, and R. Maury-Brachet. 1991. *Hexagenia rigida* (Ephemeroptera) as a biological model in aquatic ecotoxicology: experimental studies on mercury transfers from sediment. *Environmental Pollution* 69:51-67.
- Saroff, S. T. 1990. *Proceedings of the Onondaga Lake Remediation Conference*. Bolton Landing, New York: New York State Department of Law and New York State Department of Environmental Conservation. 193 pp.

Sastry, K. V. and K. Sharma. 1980. Effects of mercuric chloride on the activities of brain enzymes in a freshwater teleost, *Ophiocephalus (Channa) punctatus*. *Arch. Environ. Contam. Toxicol.* 9:425-430.

Scherer, E., F. A. J. Armstrong, and S. H. Nowak. 1975. Effects of mercury-contaminated diet upon walleyes *Stizostedion vitreum vitreum* (Mitchell). *Fish. Mar. Serv. Tech. Rep. No. 597*. Winnipeg, Manitoba: Fisheries Marine Service. 21 pp.

Schindler, D.W., S.E. Bayley, P.J. Curtis, B.R. Parker, M.P. Stanton and C.A. Kelly. 1992. Natural and man-caused factors affecting the abundance and cycling of dissolved organic substances in precambrian shield lakes. *Hydrobiologia* 229:1-21.

Schintu, M., F. Jean-Caurant, and J. C. Amiard. 1992. Organomercury determination in biological reference material: application to a study on mercury speciation in marine mammals off the Faröe Islands. *Ecotoxicology and Environmental Safety* 24:95-101.

Sharp, J. R. and J. M. Neff. 1980. Effects of the duration of exposure to mercuric chloride on the embryogenesis of the estuarine teleost, *Fundulus heteroclitus*, *Mar. Environ. Res.* 3: 195-213.

Shoichi, O. and S. Sokichi. 1985. The 1-octanol/water partition coefficient of mercury. *Bull. Chem. Soc. Japan.* 58:3401-3402.

Slooff, W., P.F.H. Bont, M. van Ewijk, and J.A. Janus. 1991. Exploratory report mercury. Report no. 710401006. Bilthoven, The Netherlands: National Institute of Public Health and Environmental Protection .

Snarski, V. M. and G. F. Olson. 1982. Chronic toxicity and bioaccumulation of mercuric chloride in the fathead minnow (*Pimephales promelas*). *Aquatic Toxicology* 2:143-156.

Spry, D. J. 1991. Metal bioavailability and toxicity to fish in low-alkalinity lakes: a critical review. *Environ. Pollut.* 71:243-304.

Spry, D. J. and J. G. Wiener. 1991. Metal bioavailability and toxicity to fish in low-alkalinity lakes: A critical review. *Environmental Pollution* 71:243-304.

- St. Louis, V.L., J.W.M. Rudd, C.A. Kelly, K.G. Beaty, N.S. Bloom and R.J. Flett. 1994. Importance of wetlands as sources of methyl mercury to boreal forest ecosystems. *Can. J. Fish. Aquat. Sci.* 51:1065-1076.
- Stromgren, T. 1982. Effect of heavy metals (Zn, Hg, Cu, Cd, Pb, Ni) on the length growth of *Mytilus edulis*. *Mar. Biol.* 72:69-72.
- Surma-Aho, K. and J. Paasivirta. 1986. Organic and inorganic mercury in the food chain of some lakes and reservoirs in Finland. *Chemosphere* 15(3):353-372.
- Swartz, R. C., P. F. Kemp, D. W. Schults, and J. O. Lamberson. 1988. Effects of mixtures of sediment contaminants on the marine infaunal amphipod, *Rhepoxynius abronius*. *Environmental Toxicology and Chemistry* 7:1013-1020.
- Syversen, T. 1977. Effects of methylmercury on in vivo protein synthesis in isolated cerebral and cerebellar neurons. *Neuropathol. Appl. Neurobiol.* 3:225-236.
- Szefer, P., W. Czarnowski, J. Pempkowiak, and E. Holm. 1993. Mercury and major essential elements in seals, penguins, and other representative fauna of the Antarctic. *Arch. Environ. Contam. Toxicol.* 25:422-427.
- Tessier, L., G. Vaillancourt, and L. Pazdernik. 1994. Temperature effects on cadmium and mercury kinetics in freshwater molluscs under laboratory conditions. *Arch. Environ. Contam. Toxicol.* 26:179-184.
- Tetra Tech, Inc. 1988. Health risk assessment of chemical contamination in Puget Sound seafood. Seattle: Environmental Protection Agency, Region 10, Office of Puget Sound. 102 pp + appendices.
- Thain, J. E. 1984. Effects of mercury on the prosobranch mollusc *Crepidula fornicata*: Acute lethal toxicity and effects on growth and reproduction of chronic exposure. *Mar. Envir. Res.* 12:285-309.
- Thompson, D. R. 1990. Metal levels in marine vertebrates. In: W. D. Furness and P. S. Rainbow (Eds.), *Heavy Metals in the Marine Environment*. pp. 144-182. Boca Raton, Florida: CRC Press.

U.S. Environmental Protection Agency (EPA). 1985. Ambient water quality criteria for mercury - 1984. U.S. EPA 440/5-84-026. Washington, D.C.: Office of Water. 136 pp.

U.S. Environmental Protection Agency (EPA). 1996. Update: National listing of fish and wildlife consumption advisories. EPA Fact Sheet EPA-823-F-96-006. Washington, D.C.: Office of Water.

U.S. Food and Drug Administration (FDA). 1984. Action level for methyl mercury in fish. Federal Register 49:45663- . November 19, 1984.

Wagemann, R., N. B. Snow, A. Lutz, and D. P. Scott. 1983. Heavy metals in tissues and organs of the narwhal (*Monodon monoceros*). Can. J. Fish. Aquat. Sci. 40(Suppl. 2):206-214.

Weber, J. H.. 1993. Review of possible paths for abiotic methylation of mercury (II) in the aquatic environment. Chemosphere 26(11):2063-2077.

Weis, J. S. and P. Weis. 1977. Effects of heavy metals on development of the killifish, *Fundulus heteroclitus*. J. Fish. Biol. 11:49-54.

Weis, P., and J.S. Weis. 1978. Methylmercury inhibition of fin regeneration in fishes and its interaction with salinity and cadmium. Estuarine Coastal Mar. Sci. 6:327-334.

Weis, J. S. and P. Weis. 1984. A rapid change in methylmercury tolerance in a population of killifish, *Fundulus heteroclitus*, from a golf course pond. Marine Environmental Research 13:231-245.

Weis, J. S., and P. Weis. 1989. Effects of environmental pollutants on early fish development. Rev. Aquat. Sci. 1:45-73.

Weis, J. S., P. Weis, and M. Heber. 1982. Variation in response to methylmercury by killifish (*Fundulus heteroclitus*) embryos. In: J.G. Pearson, R.B. Foster, and W.E. Bishop (eds.), Aquatic Toxicology and Hazard Assessment: Fifth Conference. ASTM STP 766. pp. 109-119. Philadelphia: American Society for Testing and Materials.

- Weis, J. S., P. Weis, M. Heber, and S. Vaidya. 1981. Methylmercury tolerance of killifish (*Fundulus heteroclitus*) embryos from a polluted vs non-polluted environment. *Marine Biology* 65:283-287.
- Weis, P. , J. S. Weis, and J. Bogden. 1986. Effects of environmental factors on release of mercury from Berry's Creek (New Jersey) sediments and its uptake by Killifish *Fundulus heteroclitus*. *Environmental Pollution* 40:303-315.
- Weis, P. 1984. Metallothionein and mercury tolerance in the killifish, *Fundulus heteroclitus*. *Marine Environmental Research* 14:153-166.
- Whitney, S. D. 1991. Effects of maternally-transmitted mercury on the hatching success, survival, growth, and behavior of embryo and larval walleye (*Stizostedion vitreum vitreum*). Masters Thesis. La Crosse, Wisconsin: University of Wisconsin. 71 pp.
- Wiener, J. G. and D. J. Spry. 1996. Toxicological significance of mercury in freshwater fish. In: Beyer, W.N., G.H. Heinz, and A.W. Redmon-Norwood (eds.). *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations*. Special Publication of the Society of Environmental Toxicology and Chemistry. Boca Raton, Florida: Lewis Publishers. 494 pp.
- Windom, H.L. and D. R. Kendall. 1979. Accumulation and biotransformation of mercury in coastal and marine biota. In: J.O. Nriagu (ed.) *The Biogeochemistry of Mercury in the Environment*. pp 277-302. Elsevier/North-Holland Biomedical Press 1979.
- Winfrey, M. R. and J. W. Rudd. 1990. Environmental factors affecting the formation of methylmercury in low pH lakes. *Environ. Toxicol. Chem.* 9:853-869.
- Wobeser, G. 1975. Prolonged oral administration of methyl mercury chloride to rainbow trout (*Salmo gairdneri*) fingerlings. *J. Fish.Res.Board Can.* 32:2015-2023.
- World Health Organization (WHO). 1989. *Mercury – Environmental Aspects*. International Programme on Chemical Safety (IPCS). Geneva, Switzerland. 115 pp.
- Wren, C. D. 1986. A Review of Metal Accumulation and Toxicity in Wild Mammals, I: Mercury. *Environmental Research* 40:210 -244.

Xun, L., N.E.R. Campbell, and J.W.M. Rudd. 1987. Measurements of specific rates of net methyl mercury production in the water column and surface sediments of acidified and circumneutral lakes. *Can. J. Fish. Aquat. Sci.* 44:750-757.

Zillioux, E. J., D. B. Porcella, and J. M. Benoit. 1993. Mercury cycling and effects in freshwater wetland ecosystems. *Environmental Toxicology and Chemistry* 12:1-20.